University of North Texas Health Science Center UNTHSC Scholarly Repository

Theses and Dissertations

12-1-2008

Cardiovascular Fitness and Lung Function of Adult Men and Women in the United States: NHANES 1999-2002.

Hannah Jackson *UNTHSC,* hjackson@hsc.unt.edu

Follow this and additional works at: http://digitalcommons.hsc.unt.edu/theses

Recommended Citation

Jackson, H., "Cardiovascular Fitness and Lung Function of Adult Men and Women in the United States: NHANES 1999-2002." Fort Worth, Tx: University of North Texas Health Science Center; (2008). http://digitalcommons.hsc.unt.edu/theses/3

This Thesis is brought to you for free and open access by UNTHSC Scholarly Repository. It has been accepted for inclusion in Theses and Dissertations by an authorized administrator of UNTHSC Scholarly Repository. For more information, please contact Danelle.Orange@unthsc.edu.

Jackson, Hannah L., <u>Cardiovascular Fitness and Lung Function of Adult Men and Women in the United States: NHANES 1999-2002</u>. Master of Public Health (Social & Behavioral Sciences), December 2008, 30 pp., 5 tables, 1 figure, bibliography, 70.

There is a distinct disparity between adult males and females in lung function and cardiovascular fitness in the United States. This study utilizes a nationally representative sample in order to determine predictors of lung function between men and women.

Simple means analysis, logistic and linear regressions were utilized in order to determine predictors of lung function between genders. Continuous analyses of lung function reveal that sex and BMI are the most important predictors of VO₂ max. However, analyses of clinical cut-points of cardiovascular fitness indicate that gender was not a significant predictor.

CARDIOVASCULAR FITNESS AND LUNG FUNCTION OF ADULT MEN AND WOMEN IN THE UNITED STATES: NHANES 1999-2002

Hannah L. Jackson, B.S.

APPROVED:	
Major Professor	
Committee Member	
Committee Member	
Department Chair	

Dean, School of Public Health

CARDIOVASCULAR FITNESS AND LUNG FUNCTION OF ADULT MEN AND WOMEN IN THE UNITED STATES: NHANES 1999-2002

THESIS

Presented to the School of Public Health

University of North Texas

Health Science Center at Fort Worth

in Partial Fulfillment of the Requirements

for the Degree of

Master of Public Health

By

Hannah Jackson, B.S.

Fort Worth, Texas

December 2008

Copyright by

Hannah L. Jackson

2008

ACKNOWLEDGEMENTS

I would like to thank Dr. Jim Stimpson for his guidance and mentorship throughout this thesis project. Thank you to Dr. Carlos Reyes-Ortiz and Dr. Yu-Sheng Lin as well for their encouragement, advice, helpful comments, and time spent on this project. I appreciate their willingness to serve on my thesis committee.

I would like to dedicate my thesis to my parents, Richard and Ida Jackson. The strength and perseverance they have shown me in their own lives inspired me to continue towards meeting my own academic and personal goals. I hope that in all of my success, they see their own.

I would also like to acknowledge Bradford Jackson for additional help in the statistical analysis phase of this project; and to all of my friends and family for their love and support.

H.L.J.

TABLE OF CONTENTS

		Page
LIST OF		
		vi
LIST OF		
FIGURES		V11
Chapter		
1. INTROD	DUCTION	1
2. LITERA	ATURE REVIEW	2
Life	fe Span Epidemiology of Lung Function	
	cioeconomic Determinants of Lung Function	
Hea	ealth Behaviors	
Sta	atement of Problem	
3. METHO	ODOLOGY	11
Dat	uta	
Der	ependent Variable	
Co	ovariates	
Sta	atistical Analysis	
4. RESULTS		13
5. DISCUSSION	ON AND CONCLUSIONS	16
REFERENCES		19

LIST OF TABLES

1.	Sample Characteristics Stratified by Sex: NHANES 1999-	
	2002	25
2.	Weighted Percentages of Cardiovascular Fitness	
_	Level	26
	Weighted Means (Standard Errors) of VO ₂ max	
		27
4.	Linear Regression of VO ₂ max on Selected Independent	
	Variables	28
5.	Logistic Regression of High Cardiovascular Fitness Independent	
	Variables	29

LIST OF FIGURES

1.	Determinants of Lung Function Conceptual	
	Model	.30

CHAPTER ONE

INTRODUCTION

Over 35 million Americans suffer from chronic lung diseases such as cancer, asthma, chronic obstructive pulmonary disease (COPD), and bronchitis ¹. Respiratory diseases are the fourth leading cause of death in the United States, preceded only by heart disease, cancers, and cerebrovascular disease ². Growing levels of lung disease coupled with declining levels of lung function are global health issues that are modifiable and preventable.

Respiratory diseases are a leading cause of death in women. In the past year, more women died of COPD and chronic bronchitis than men and lung cancer has become the leading cancer cause of death in women ¹. Furthermore, other chronic lung diseases have increased more rapidly for women than men. For example, in the past decade, asthma rates have risen by 97% in women in comparison to a 22% increase for men.

The reason for increasing differences in lung function between men and women has not been studied. The purpose of this study is to assess and explain differences in lung function between men and women using national data. It is expected that changes in social and behavioral predictors of lung function such as environmental exposure, socioeconomic status, access to care, and health behaviors have contributed to differences in lung function between men and women. The public health importance of these findings are the potential identification of modifiable mechanisms to reduce or eliminate sex differences in lung function and improve lung health for the US population.

CHAPTER TWO

LITERATURE REVIEW

Life Span Epidemiology of Lung Function

Throughout the human lifespan, female lungs are consistently smaller in size and in weight compared to male lungs. At birth, the female lungs have fewer respiratory bronchioles resulting in restricted airflow to the interior of the lungs³. During the first few weeks of life, significant changes and differences in lung function and development become apparent between men and women ⁴. Significantly larger tidal flow volume differences are apparent in infant boys as early as the first few days of life ⁵. Lower airway function in early infancy has serious implications for increased susceptibility to lung disease in childhood and serves as a risk factor for airflow obstruction in adulthood⁶. Despite lower levels of lung function at early stages of life, infant girls exhibit a more rapid maturation of lungs the first few weeks of life compared with boys ⁷.

Throughout the lifespan, female lung growth occurs more rapidly than males but for a shorter duration. Maximum lung growth is complete in females early in adolescence while male lungs continue to develop into early adult life ⁸, which is associated with increased lung capacity versus male counterparts at this stage of life ⁹. Lung development in females peaks early in the adolescence years, whereas for males, thoracic growth continues into early adulthood ¹⁰. Unlike with females, pulmonary function measurements in males indicate continued lung growth and development with age into early adulthood ¹¹.

Adult female lungs are consistently smaller in capacity and weight, and consequently contain fewer respiratory bronchioles ¹². Another contributing factor for greater lung function in men is the increase in width of the thorax and trachea where women do not demonstrate any change ¹³. Height-matched men have larger diameter airways and larger lung volumes and diffusion surfaces ¹⁴. Age related loss of lung function occurs in both genders; however healthy males exhibit higher levels of lung function, yet women experience greater rates of decline in respiratory function ¹⁵. In older adult females, lung function begins to rapidly decline at an earlier age and at a faster rate than age-matched male counterparts ¹⁶. Although early life events have a strong influence on lung function between men and women, the growth process may only partially account for sex differences in lung function.

Socioeconomic Determinants of Lung Function

Socioeconomic factors have a significant impact on lung performance and vulnerability to lung disease. Between genders, social influences are important determinants in lung health outcomes. Social factors and roles such as occupational status, education and income level, as well as utilization of health care services that differ between men and women may partially explain disparities in lung function between the genders.

Environmental and occupational exposures have a strong influence on lung function and health. Coarse and fine airborne particles and various pollutants contribute to an increased risk of respiratory damage and disease. There is a high correlation

between fine particle concentration and asthma related response that include lower respiratory symptoms and decreased peak flow ¹⁷. Exposure to ambient concentrations of ozone is related to lower levels of lung function and significant reductions in airway size ¹⁸.

Women experience greater respiratory effects of ambient particles and air pollution exposure compared to men. There is a greater mortality rate observed in women than men due to ambient particles. Particles deposit in all regions of lung differently between genders ¹⁹. In women, there is suggestive evidence that there is an increased susceptibility with exposure to particulate matter in inflammatory response of the respiratory system ²⁰.

Living in urban areas in proximity to traffic dense areas and industrial air pollution are also primary environmental risk factors for respiratory diseases. Rural communities are far removed from industrial air pollution, automobile traffic emissions, and other air quality and health that individuals in urban areas address. Residents of rural areas report overall better health and health behavior practices compared with their urban counterparts ²¹.

Respiratory health and chronic respiratory conditions have a lower prevalence in rural communities than urban regions, and it has been suggested that the rural environment serves as a protective factor against developing chronic lung diseases²².

Asthma prevalence rates are consistently lower among farm children compared to non-farm children. Differences in environmental exposures partially explain this phenomenon

in addition to agricultural and livestock exposures that serve as a protective component in the rural environment against developing various lung diseases ²³. Although rural children have better overall lung function than urban counterparts, rural girls exhibit a higher percentage of wheezing, asthma and airway hyper responsiveness than rural boys²⁴.

Alternatively, urban area residents experience higher prevalence rates of respiratory diseases as a result of environmental hazards that are present in their environment. Urban area neighborhoods have a closer proximity to industrial and traffic pollution that are major risk factors for developing airway and lung disease ²⁵. Traffic exposure is a primary risk factor in urban areas for developing various lung diseases and lowering lung function. Area census tracts that face intersections and highways have a 60 % increased risk of developing asthma compared to areas located away from major road sites ²⁶. Fine particulates are 29% more dense in the air in these areas compared with residential areas located further away ²⁷. There is no significant effect of traffic density on lung function in men; however women experience significant FEV₁ and FVC decline with higher traffic density exposure compared to subjects living further away from traffic dense areas ²⁸.

Occupational exposure to vapors, dust, or fumes is more common among males than among females. Trends indicate that men occupy a majority of the manual labor and construction fields. These occupational environments increase the amount of exposure and susceptibility to dust, mold, and various outdoor pollutants that may increase the risk

for adverse lung conditions ²⁹. However, among women, occupational status showed a significant difference in lung function outcomes. Women who hold administrative and higher socioeconomic positions showed decreased odds of developing chronic lung diseases compared to women in industrial and agricultural positions ³⁰. Because women are less likely to hold jobs that expose them to various ambient particulate matter, better lung function is observed in occupational field compared to men.

Lung function is positively associated with educational achievement and income status. This relationship is consistent with other socioeconomic factors where higher levels of education and income demonstrate better overall lung function than individuals with lower education and income status ³¹. Maternal education is a strong predictor of lung function in children. Children born to mothers with lower levels of have lower FEV1 values compared to children whose maternal education consisted of at least a college degree ³².

Income level and educational status are closely correlated with one another. Household income levels exhibit a positive correlation with lung function and general health; where higher household income levels yield higher FEV₁ and FVC values in both women and men compared to lower income families ³³. Both education and income levels are important determinants of lung function in men and women. These factors are strong predictors of optimal lung and overall health.

Health care access and utilization differences in gender and symptomatic individuals have a significant impact on health policy and management in the United States. Perceptions and attitudes towards illness and disease can act as an important

determinant in diagnosis and detection of lung disease ³⁴. Patients with severe lung disease such as COPD use hospital care 2.3 times more than age and gender matched asymptomatic individuals. Accurate and early diagnoses of lung disease are important determinants of lung health³⁵. Standards for treatment and detection of lung diseases have improved with the use of spirometric techniques for diagnosing chronic lung diseases. Spirometric data are also used to determine individual lung capacity and volume. Surveys of chest physicians indicated supplementary use of chest x-rays and spirometry in addition to standard clinical data and reported symptoms in order to accurately diagnosis lung disease in their patients ³⁶. However, a distinct bias exists between genders in the diagnosis of chronic lung diseases in women. When compared to men, women are consistently under-diagnosed with COPD and other chronic lung conditions despite similar presented symptoms ³⁷. In initial diagnosis and treatment of lung diseases, there are significant gender differences in the attitudes of physicians in diagnosing women with lung disease. Despite lower pack-years of smoking and increased reported symptoms of respiratory difficulties, physicians failed to order supplemental testing for females that resulted in an overall under-diagnosis of COPD ³⁸. This disparity in diagnoses can partially account for poor lung health among women.

Health Behaviors

Health disparities between men and women are fundamentally associated with behavioral practices and lifestyle decisions. In general women lead healthier lifestyles and engage in healthy behaviors more often than men, yet demonstrate lower levels of lung function and overall health. Occupational roles and status of men compared to women in part explains this gender health paradox ³⁹.

Tobacco exposure is one of the major risk factors in the development of lung disease and is responsible for the rapid decline in lung function. Tobacco smoking plays a major role in the mortality of U.S. women, where 90% of all lung cancer deaths among women are attributable to smoking 40. One third of the male population in the world smokes, while the number of women who are exposed to tobacco is greater than previous generations and continues to 41. Smoking during pregnancy is responsible for reduced lung function in infants as early as the first 4-5 weeks of life. Parental smoking is associated with an increased number of lower respiratory tract infections, and the presence of persistent cough and phlegm in children of smoking parents 41. A clear interaction between chronic bronchitis and airflow obstruction is observed in addition to decline in forced expiratory volume in one second (FEV₁) amongst ever-smokers 42.

While it is known that smoking is associated with reduced lung function in general, women experience a greater overall loss of lung function than men ⁴³. Although smoking habits in women are less prevalent than in men, females experience a higher proportion of respiratory symptoms compared to males despite a lower prevalence of smoking status ⁴⁴. In the effects of cigarette smoking, women with COPD demonstrate greater levels of breathlessness relative to men at similar degrees of airflow obstruction and emphysema severity ⁴⁵. Likewise, women report smoking fewer pack-years than men but demonstrate similar levels of COPD severity as men ⁴⁶. In women with no history of smoking, passive tobacco exposure is shown to moderately reduce lung

function, whereas in women with asthma reduction in lung function is significantly greater than that of the general population ⁴⁷.

Participation in exercise is positively associated with higher levels of lung function both in healthy, non-smoking males and females. However, non-smoking men demonstrated higher FEV₁, FVC levels after exercise than non-smoking women ⁴⁸. Bronchial hyper responsiveness, a characteristic of asthma and inflammation of the airways, is negatively associated with increased frequency and duration of physical activity. In asymptomatic women, after adjusting for confounding factors, FEV₁ levels are still consistently lower in women after exercise than age-matched men ⁴⁹. While exercise poses as a protective factor in expanding the lung and increasing respiratory function, its overall effect on individual health is minimal.

Nutrition and dietary intake are both associated with the cause and prevention of various diseases. Through cell development, regulation of the airway smooth muscle, and enzymatic reaction of neuromuscular transmission, an adequate diet containing a balance of omega 3 fatty acids, antioxidants and electrolytes; may inhibit various reactions and promote cellular activities in the lungs that reduce an individual's susceptibility to adverse lung health ⁵⁰. In children, an adequate diet consisting of recommended daily allowances of nutrients reflects a greater effect on airflow in girls than boys ⁵¹. This phenomenon could reflect the early anatomical developmental differences between genders. Limited research in the Nurses' Health Study has shown that higher levels of antioxidants and vitamins in dietary consumption act as a protective factor against COPD development in female nurses ⁵².

Statement of Problem

At present, what is not known are variables that effect cardiovascular fitness and lung function between men and women. Previous research has focused on adolescent cardiovascular fitness and disparities between race/ethnicities. This study will use nationally representative data to identify sex differences in lung function. It is expected that women will have lower lung function than men. Further, social and behavioral factors are expected to explain the majority of the difference in lung function between men and women. Therefore, the findings from this study could be used to build effective intervention efforts to improve lung function for the United States population and reduce disparities in lung function between men and women.

CHAPTER THREE

METHODOLOGY

Data

This study uses data from the 1999-2000, and 2001-2002 National Health and Nutrition Examination Surveys. These data are publicly available for download from the National Center for Health Statistics, which is part of the Center for Disease Control and Prevention. NHANES is a stratified, multi-staged monitoring survey that uses probability sampling to evaluate the health and nutritional status of the civilian, non-institutionalized population of the United States. Further details about NHANES can be found elsewhere 53-54. Individuals will be eligible if they are female or male, aged 20-49 years, and provided a reproducible VO₂ max score.

Dependent Variable

Cardiovascular fitness is measured by conducting exercise stress tests on a treadmill or cycle ergometer through VO₂ max values. VO₂ max is the maximum amount of oxygen uptake, measured in liters of oxygen per minute (l/min) or milliliters of oxygen per kilogram of weight per minute (ml/kg/min), and is considered the best indicator of cardiovascular health and lung function⁵⁵. Measurement of VO₂ max involves a physical effort in duration and intensity to fully tax the aerobic system. In clinical testing, involves a graded exercise test (either on a treadmill or on a cycle ergometer) in which exercise intensity is incrementally increased while measuring ventilation of oxygen and carbon dioxide concentration of the inhaled and exhaled air⁵⁶. VO₂ max is reached when oxygen

consumption remains at steady state despite an increase in workload. During the process, the amounts of oxygen utilized and carbon dioxide expired are measured at a predetermined sub maximal exercise capacity to determine the VO_2 max⁵⁷⁻⁵⁹.

Using sex and age-specific criteria from NHANES data, the estimated VO₂ max will be categorized as a low, moderate or high level of cardiovascular fitness and lung function outcome based on data from the Aerobic Center Longitudinal Study a VO₂ max below the 20th percentile of the Aerobic Center Longitudinal Study; a moderate fitness level will be a value between the 20th and 59th percentile, and high cardiovascular fitness will be defined as above the 60th percentile.⁶⁰

Covariates

Socioeconomic indicators include age, race/ethnicity, and education. Health behaviors include smoking status, exercise, and diet. Anatomical control variables will be assessed by body mass index (height in meters /weight in kilograms-squared).

Statistical Analysis

A simple means analysis will be used to observe the differences in lung function between groups. Results will be stratified by age and sex. Multivariate linear and logistic regression models will be used to adjust for factors predictive of lung function if these factors explain the differences in lung function for men and women. All analyses will be adjusted for sample weights, strata, and population sampling unites using STATA 10.0 (StatCorp, College Station, TX). P-values less than 0.05 will be considered statistically significant.

CHAPTER FOUR

RESULTS

Description of the Sample

Table 1 describes the variables used in the study stratified by sex. Given that the National Health and Nutrition Examination Survey is a nationally representative sample of the United States population, the demographic characteristics of the sample closely match figures from the US Census. Overall, 51% of the sample was female and 48.6 were male. The race/ethnic distribution was 38% Non Hispanic White and 62% non-White which includes non Hispanic Black, Mexican-Americans, Hispanics and Asians. Over half of the sample has a high school degree (58%). With regard to health behaviors, compared to males, females had higher body mass index, higher serum triglyceride, lower participation moderate activity, and lower serum cotinine. The outcome is cardiovascular fitness level based upon sex-age cut points of estimated VO₂ max. In this sample, women had a lower level of VO₂ max (37.92) compared to men at 45.85. The average VO₂ max values for this cross sectional sample were 42.0. Males had a higher proportion of high cardiovascular fitness (35%) compared to women (32%) and women had a higher proportion of low cardiovascular fitness (29%) than men (26%).

Association of Sex and Cardiovascular Fitness by Age

Table 2 presents the weighted percentage of cardiovascular fitness levels for males and females by age. Among persons 20-29 years of age, there were no significant

difference between men and women for low cardiovascular fitness (13%). However more men 20-29 (23%) achieved high levels of cardiovascular fitness than women (20%). Women 30-39 years of age had a higher percentage (12%) of low cardiovascular fitness than men (11%). Women 40-49 years of age had a higher percentage among low cardiovascular fitness levels (5%) compared with male counterparts (4%). Overall average indicates that women had higher percentages (10%) of low cardiovascular fitness levels than men (9%). Men also had a higher total percentage (23%) of high cardiovascular fitness level compared to 20% of overall women.

Table 3 shows the weighted average VO₂ max values for males and females stratified by age group. Averages in the overall categories indicate that men across all age groups had, on average, higher VO₂ max values (45.02) than females (36.82). Consistently across all age groups men had significantly higher mean VO₂ max values than women. Men between the ages of 20-29 had an average VO₂ max value of 46.07 compared to women counterparts at 37.96. Men in the 30-39 age range in this sample had a VO₂ max average of 45.33 compared to 37.32 for women. In the highest age bracket, 40-49 years, men continued to show high VO₂ max values (43.92) and women significantly lower (35.20).

Multivariate Association of Sex and Cardiovascular Fitness

Table 4 provides standardized and unstandardized coefficients from the linear regression of VO₂ max with adjustment for the complex sampling design of NHANES. Overall, the regression model explained 18% of the variance for VO₂max. All

independent variables are statistically significant except for serum cotinine and physical activity. White race was positively associated with VO_2 max. All other variables were negatively associated with VO_2 max. Standardized coefficients are provided to compare which predictor variables are most strongly associated with VO_2 max. Sex is substantially larger than all other standardized coefficients suggesting that it may be the most important predictor of VO_2 max in this analysis. Body mass index was the next strongest predictor followed by education, serum triglycerides, age, serum cotinine, physical activity, and race.

Table 5 presents results from logistic regression analysis of high cardiovascular fitness with adjustment for the complex sampling design of NHANES. In this analysis, only age, race, and body mass index were statistically significant predictors of high cardiovascular fitness. All other variables were not statistically significant. Other factors equal, high cardiovascular fitness level was positively associated with age and white race and negatively associated with body mass index. These results suggest that, other factors equal, white race, more years of age are associated with a higher likelihood of being fit, while higher levels of body mass index are associated with a lower likelihood of being fit. Unlike in the linear regression results where sex was found to be the most important predictor of VO_2 max, in the logistic regression analysis sex was not found to be statistically significant.

CHAPTER 5

DISCUSSION AND CONCLUSIONS

This study sought to identify behavioral and socioeconomic determinants of lung function between men and women in the United States using a nationally representative sample. Cardiovascular fitness level cut-points and VO₂ max assessments were used to measure lung function. Overall, there are important gender differences in lung function in this nationally representative sample of adults. Gender was the strongest predictor of lung function, where females had lower levels of VO₂ max and cardiovascular fitness consistently across all age groups. In addition to gender, key predictors of lung function were body mass index, race/ethnicity, education and age.

Analysis of the continuous measure of lung function revealed that sex and body mass index were the two most important predictors of VO_2 max. However, analysis of the clinical cut-point measure suggested that race/ethnic minority status and age were associated with better lung function, and gender was not a significant predictor. Taken together, these results suggest a potential problem with the clinical cardiovascular fitness cut-points. Age and minority status should be negatively associated with lung function as has been found in other studies using alternative measurement devices such as spirometry $^{61-67}$.

Furthermore, given that gender was the strongest predictor of the continuous measure of lung function, one would expect that gender should at least be among the significant predictors in the binary categorization of lung function. Therefore, this study

provides support for a previous study that argued continuous estimates of VO_2 max levels more accurately represent lung function as opposed to age-sex categorization of cardiovascular fitness⁶⁸.

In the context of current lung function literature regarding lung function and cardiovascular fitness, this study highlights the disparities in predictor variables of cardiovascular fitness levels. Clinical cardiovascular fitness cut-points established by the Aerobics Center Longitudinal Study (ACLS) classify cardiovascular fitness into low, medium and high categories based upon estimated VO₂ max values. However, results from this study suggest that more investigation is needed into the classification of cardiovascular fitness in these clinical categorizations due to the differences in predictors of VO₂ max.

Strengths and Limitations

The principal strength of this study is the use of a large, nationally representative sample with clinical assessment for the outcome measure and several control variables. However, this study was limited by the cross-sectional nature of the study design which precluded causal inferences. Another limitation is that some measures such as physical activity were self-reported. Results should also be reviewed cautiously because of the procedures used to assess cardiovascular fitness. Exclusion criterion for participation in VO₂ max and cardiovascular fitness testing of the NHANES included age, physical and other medical and physical limitations ⁶⁹⁻⁷⁰. Therefore there is the potential for selection bias of a healthier sample compared to the broader population. Finally, this study was

unable to include other important determinants of lung functions such as exposure to toxins.

Conclusion

Given that the resources for public health initiatives are increasingly scarce, and that lung health is an under-promoted area of public health research, future research is needed in this area to further investigate differences in gender and lung function. Future direction in this area should focus on examining trends in lung function using cardiovascular fitness over an extended time period to accurately assess patterns and predictors of respiratory health in the United States. Furthermore, this study indicates a disparity between clinical cardiovascular fitness categorizations and a VO₂ max values, which indicate a need for more research into lung function disparities.

REFERENCES

- 1. American Lung Association. *Data and Statistics*. Available at: http://www.lungusa.org/site/c.dvLUK9O0E/b.33347/2008.
- 2. Centers for Disease Control and Prevention, National Center for Health Statistics. Deaths- Leading Causes .Hyattsville, MD: US Department of Health and Human Services, Centers for Disease Control and Prevention. Internet: http://www.cdc.gov/nchs/fastats/lcod.htm (accessed July 2008)
- 3. Marieb,EM. "Human Anatomy and Physiology, 6th Edition". Pearson Education Inc. San Fransico, 2004.p. 835-37
- 4. Friedrich L, Stein RT, Pitrez PM, Corso AL, Jones MH. Reduced lung function in healthy preterm infants in the first months of life. *Am J Respir Crit Care Med*. 2006;173:442-7.
- 5. Lødrup Carlsen KC, Magnus P, Carlsen KH. Lung function by tidal breathing in awake healthy newborn infants. *Eur Respir Journal*. 1994;7:1660-8.
- 6. Stern DA, Morgan WJ, Wright AL, Guerra S, Martinez FD. Poor airway function in early infancy and lung function by age 22 years: A non-selective longitudinal cohort study. *Lancet*. 2007;370:758-764.
- 7. Becklake MR, Kauffmann F. Gender differences in airway behaviour over the human life span. *Thorax*. 1999;54:1119-1138.
- 8. Carey MA, Card JW, Voltz JW, et al. It's all about sex: Gender, lung development and lung disease. *Trends Endocrinol Metab*. 2007;18:308-313.
- 9. Nielsen KG. Lung function and bronchial responsiveness in young children. clinical and research applications. *Dan Med Bull*. 2006;53:46-75.
- 10. Varkey AB. Chronic obstructive pulmonary disease in women: Exploring gender differences. *Curr Opin Pulm Med.* 2004;10:98-103.
- 11. Nève V. Lung and thorax development during adolescence: Relationship with pubertal status. *Eur Respir Journal*.. 2002;20:1292-8.
- 12. Carey MA, Card JW, Voltz JW, Germolec DR, Korach KS, Zeldin DC. The impact of sex and sex hormones on lung physiology and disease: Lessons from animal studies. *Am J Physiol Lung Cell Mol Physiol*. 2007;293:L272-8.
- 13. Boezen HM. Sex and gender differences in lung development and their clinical significance. *Clin Chest Med.* 2004;25:237-45.

- 14. Harms CA. Does gender affect pulmonary function and exercise capacity? *Respir Physiol Neurobiol*. 2006;151:124-131.
- 15. Vollmer WM. Patterns of lung function in asymptomatic nonsmoking men and women in the people's republic of china. *Ann Epidemiol*. 2002;12:295-302.
- 16. Griffith KA. Predictors of loss of lung function in the elderly: The cardiovascular health study. *Am J Respir Crit Care Med*. 2001;163:61-8.
- 17. Schwartz J. Fine particles are more strongly associated than coarse particles with acute respiratory health effects in schoolchildren. *Epidemiology*. 2000;11:6-10.
- 18. Tager IB, Balmes J, Lurmann F, Ngo L, Alcorn S, Kunzli N. Chronic exposure to ambient ozone and lung function in young adults. *Epidemiology*. 2005;16:751-759.
- 19. Zanobetti A. Race, gender, and social status as modifiers of the effects of PM10 on mortality. *J Occup Environ Med.* 2000;42:469-74.
- 20.. Franklin M, Zeka A, Schwartz J. Association between PM2.5 and all-cause and specific-cause mortality in 27 US communities. *J Expo Sci Environ Epidemiol*. 2007;17:279-287.
- 21. Iversen L, Hannaford PC, Price DB, Godden DJ. Is living in a rural area good for your respiratory health? results from a cross-sectional study in scotland. *Chest*. 2005;128:2059-2067.
- 22. Sigsgaard, T. Respiratory Health and Allergy Among Young Farmers and Non-Farming Rural Males in Denmark: The *SUS* Study. *Journal of Agromedicine* 1997: 4(12): 63-78.
- 23. Naleway AL. Asthma and atopy in rural children: Is farming protective? *Clin Med Res*. 2004;2:5-12.
- 24. Ernst P, Cormier Y. Relative scarcity of asthma and atopy among rural adolescents raised on a farm. *Am J Respir Crit Care Med*. 2000;161:1563-1566.
- 25. Holguin F. Traffic related exposures and lung function in adults. *Thorax*. 2007;62:837-838.
- 26. Juhn YJ, Sauver JS, Katusic S, Vargas D, Weaver A, Yunginger J. The influence of neighborhood environment on the incidence of childhood asthma: A multilevel approach. *Soc Sci Med*. 2005;60:2453-2464.

- 27. Yang HH, Hsieh LT, Lin MC, Mi HH, Chen PC. Dry deposition of sulfate-containing particulate at the highway intersection, coastal and suburban areas. *Chemosphere*. 2004;54:369-378.
- 28. Kan H, Heiss G, Rose KM, Whitsel E, Lurmann F, London SJ. Traffic exposure and lung function in adults: The atherosclerosis risk in communities study. *Thorax*. 2007;62:873-879.
- 29. Zock JP, Sunyer J, Kogevinas M, Kromhout H, Burney P, Anto JM. Occupation, chronic bronchitis, and lung function in young adults. an international study. *Am J Respir Crit Care Med.* 2001:163:1572-1577.
- 30. Krstev S. Occupation and chronic bronchitis among chinese women. *J Occup Environ Med*. 2008;50:64-71.
- 31. Hegewald MJ, Crapo RO. Socioeconomic status and lung function. *Chest.* 2007;132:1608-1614.
- 32. Lercher P, Schmitzberger R. Birth weight, education, environment, and lung function at school age: A community study in an alpine area. *Eur Respir J.* 1997;10:2502-2507.
- 33. Lurie N. Health disparities and access to health. *JAMA*. 2007;297:1118-21.
- 34. Merzel C. Gender differences in health care access indicators in an urban, low-income community. *Am J Public Health*. 2000;90:909-916.
- 35. Schermer T. Referral and consultation in asthma and COPD: an exploration of pulmonologists' views. *Neth J Med.* 2003: 62(3): 71-81.
- 36. Sivori ML, Raimondi GA. Survey of chest physicians regarding COPD diagnosis and treatment. *Medicina (B Aires)*. 2004;64:113-119.
- 37. Chapman KR. Gender bias in the diagnosis of COPD. Chest. 2001;119:1691-5.
- 38. Watson L, Vestbo J, Postma DS, et al. Gender differences in the management and experience of chronic obstructive pulmonary disease. *Respir Med.* 2004;98:1207-1213.
- 39. Ross CE. Sex stratification and health lifestyle: Consequences for men's and women's perceived health. *J Health Soc Behav.* 1994;35:161-78.
- 40. Surgeon General's Report- Women and Smoking; 2001. Internet: http://www.cdc.gov/tobacco/data_statistics/sgr/sgr_2001/sgr_women_chapters.htm (accessed June 2008).

- 41. Carlsen KH, Carlsen KC. Respiratory effects of tobacco smoking on infants and young children. *Paediatr Respir Rev.* 2008;9:11-9; quiz 19-20.
- 42. Pelkonen M. Smoking: Relationship to chronic bronchitis, chronic obstructive pulmonary disease and mortality. *Curr Opin Pulm Med.* 2008;14:105-109.
- 43. Prescott E, Bjerg AM, Andersen PK, Lange P, Vestbo J. Gender difference in smoking effects on lung function and risk of hospitalization for COPD: Results from a danish longitudinal population study. *Eur Respir J.* 1997;10:822-827.
- 44. Langhammer A, Johnsen R, Gulsvik A, Holmen TL, Bjermer L. Sex differences in lung vulnerability to tobacco smoking. *Eur Respir J.* 2003;21:1017-1023.
- 45. Martinez FJ, Curtis JL, Sciurba F, et al. Sex differences in severe pulmonary emphysema. *Am J Respir Crit Care Med*. 2007;176:243-252.
- 46. DeMeo DL. The yin and yang of COPD: Sex/gender differences in the national emphysema treatment trial. *Am J Respir Crit Care Med*. 2007;176:222-223.
- 47. Eisner MD. Environmental tobacco smoke exposure and pulmonary function among adults in NHANES III: Impact on the general population and adults with current asthma. *Environ Health Perspect*. 2002;110:765-770.
- 48. Holmen TL, Barrett-Connor E, Clausen J, Holmen J, Bjermer L. Physical exercise, sports, and lung function in smoking versus nonsmoking adolescents. *Eur Respir J*. 2002;19:8-15.
- 49. Shaaban R. Physical activity and bronchial hyperresponsiveness: European community respiratory health survey II. *Thorax*. 2007;62:403-10.
- 50. Romieu I. Nutrition and lung health. *Int J Tuberc Lung Dis.* 2005;9:362-74.
- 51. Gilliland FD, Berhane KT, Li YF, Kim DH, Margolis HG. Dietary magnesium, potassium, sodium, and children's lung function. *Am J Epidemiol*. 2002;155:125-131.
- 52. Varraso R. Prospective study of dietary patterns and chronic obstructive pulmonary disease among US women. *Am J Clin Nutr.* 2007;86:488-95.
- 53. Centers for Disease Control and Prevention, National Center for Health Statistics. National Health and Nutrition Examination Survey: NHANES 1999-2000 data files documentation codebook, 1999-2000. Hyattsville, MD: US Department of Health and Human Services, Centers for Disease Control and Prevention. Internet: http://www.cdc.gov/nchs/about/major/nhanes99_00.htm (accessed July 2008)

- 54. Centers for Disease Control and Prevention, National Center for Health Statistics. National Health and Nutrition Examination Survey: NHANES 2001-2002 data files documentation codebook, 2000-2001. Hyattsville, MD: US Department of Health and Human Services, Centers for Disease Control and Prevention. Internet: http://www.cdc.gov/nchs/about/major/nhanes01-02.htm (accessed July 2008
- 55. Thomas E. Hyde and Marianne S. Gengenbach, *Conservative Management of Sports Injuries* (2nd ed; Sudbury, Mass.: Jones & Bartlett, 2007), 845.
- 56. Foster C, Hector L, McDonald K, Snyder A. Measurement of anaerobic power and capacity. In: Maud P, Foster C, eds. *Physiological assessment of human fitness*. Champaign, IL.: Human kinetics, 1995:73-85.
- 57. Howley ET, Bassett DR Jr, Welch HG. Criteria for maximal oxygen uptake: review and commentary. *Med Sci Sports Exerc*. 1995 Sep 27(9):1292-301.
- 58 Duncan GE, Howley ET, Johnson BN. Applicability of VO2 max criteria: discontinuous versus continuous protocols. *Med Sci Sports Exerc*. 1997 Feb;29(2):273-8.
- 59 Poole DC, Richardson RS. Determinants of oxygen uptake. Implications for exercise testing. *Sports Med.* 1997 Nov;24(5):308-20.
- 60. Blair, S.N., H.W. Kohl, III, R.S. Paffenbarger, Jr., D.G. Clark, K.H. Cooper, L.W. Gibbons. Physical Fitness and all cause mortality: a prospective study of healthy men and women. *JAMA* 262:2395-2401,1989.
- 61. Enright PL, Kronmal RA, Higgins MW, Schenker MB, Haponik EF. Prevalence and correlates of respiratory symptoms and disease in the elderly. Cardiovascular Health Study. *Chest.* 1994 Sep;106(3):827-34.
- 62. McClaran SR, Babcock MA, Pegelow DF, Reddan WG, Dempsey JA. Longitudinal effects of aging on lung function at rest and exercise in healthy active fit elderly adults. *J Appl Physiol*. 1995 May;78(5):1957-68.
- 63. Stamford BA. Exercise and the elderly. Exerc Sport Sci Rev. 1988;16:341-79.
- 64. Haselkorn T, Lee JH, Mink DR, Weiss ST; TENOR Study Group. Racial disparities in asthma-related health outcomes in severe or difficult-to-treat asthma. *Ann Allergy Asthma Immunol.* 2008 Sep;101(3):256-63.
- 65. Whitrow MJ, Harding S. Ethnic differences in adolescent lung function: anthropometric, socioeconomic, and psychosocial factors. *Am J Respir Crit Care Med*. 2008 Jun 1;177(11):1262-7. Epub 2008 Mar 6.

- 66. Panico L, Bartley M, Marmot M, Nazroo JY, Sacker A, Kelly YJ. Ethnic variation in childhood asthma and wheezing illnesses: findings from the Millennium Cohort Study. *Int J Epidemiol*. 2007 Oct;36(5):1093-102. Epub 2007 May 24
- 67. Akinbami LJ, Rhodes JC, Lara M. Racial and ethnic differences in asthma diagnosis among children who wheeze. *Pediatrics*. 2005 May;115(5):1254-60.
- 68. 61. Powers, SK. "Exercise Physiology. Theory and Application to Fitness and Performance" 5th Edition, p.288
- 69. Centers for Disease Control and Prevention, National Center for Health Statistics. National Health and Nutrition Examination Survey: NHANES 1999-2000 data release MEC documentation, 1999-2000. Hyattsville, MD: US Department of Health and Human Services, Centers for Disease Control and Prevention. Internet: http://www.cdc.gov/nchs/data/nhanes/frequency/cvx_doc.pdf (accessed November 2008)
- 70. Centers for Disease Control and Prevention, National Center for Health Statistics. National Health and Nutrition Examination Survey: NHANES 2000-2001 data release MEC documentation, 2000-2001. Hyattsville, MD: US Department of Health and Human Services, Centers for Disease Control and Prevention. Internet: http://www.cdc.gov/nchs/data/nhanes/nhanes_01_02/cvx_b_doc.pdf (accessed November 2008

Table 1. Sample Characteristics Stratified by Sex: NHANES 1999-2002

TOTAL	MEN	WOMEN	TOTAL	MEN	WOMEN	TOTAL
	N	Mean (SE)	N	Mean (SE)	N	Mean (SE)
		or %		or %		or %
Female, %	21,004	100	10,214	48.63	10,790	51.37
Age, years	21,004	29.5 (24.9)	10,214	29.0 (0.25)	10,790	30.0 (0.24)
White, %	21,004	62	10,214	62	10,790	61.5
High School or Less,	17,498	42	8,415	40	9,083	43
%						
Body Mass Index,	17,472	25.0 (7.03)	8,478	24.35	8,994	25.21 (0.08)
Serum Triglycerides,	3,503	124(95.4)	1,725	123.41	1,778	124.35
mg/dl				(2.38)		(2.18)
Serum Cotinine,	15,583	33.4 (93.0)	7,596	42.31	7,987	25.04 (0.88)
ng/ml				(1.20)		
Physical Activity, %	14,965	44	7,142	45	7,823	44
Cardiovascular	5,515		2,822		2,693	
Fitness, %		27		26		29
Low		38		39		39
Medium		34		35		32
High						
VO ₂ max, ml/min/kg	5,515	42 (10.3)	2,822	45.85	2,693	37.92 (0.17)
				(0.19)		

To convert ng/mL cotinine to nmol/L, multiply by 5.68. To convert mg/dL triglycerides to mmol/L, multiply by 0.0112.

Table 2: Weighted Percentages of Cardiovascular Fitness Level by Age for Men and Women

	MEN					WO	MEN	
	Total	20-29	30-39	40-49	Total	20-29	30-39	40-49
Low	9%	13%	11%	4%	10%	13%	12%	5%
Medium	19%	21%	20%	15%	18%	19%	18%	16%
High	23%	19%	21%	21%	20%	14%	18%	28%
N	5515	4111	4875	640	5515	4111	4875	640

Table 3: Weighted Means (Standard Errors) of VO₂ max by Age for Men and Women

	MEN			WOMEN				
	Total	20-29	30-39	40-49	Total	20-29	30-39	40-49
Mean (SE)	45.02	46.07	45.33	43.92	36.82	37.96	37.32	35.20
	(0.28)	(0.36)	(0.31)	(0.78)	(0.29)	(0.29)	(0.30)	(0.49)
N	5515	2114	2496	326	5515	1997	2379	314

Table 4: Linear Regression of VO₂ max on Selected Independent Variables

	Standardized Coefficient	Unstandardized Coefficient	Standard Error	P- Value
Female	-0.32	-7.03	0.94	0.00
Age	-0.04	-0.04	0.03	0.30
White	0.00	1.2	0.87	0.20
Education	-0.09	-1.95	0.98	0.10
Body Mass Index	-0.14	-0.15	0.07	0.05
Serum	-0.07	-0.01	0.00	0.02
Triglycerides Serum Cotinine	0.02	-0.00	0.00	0.91
Physical Activity	-0.02	-0.24	0.71	0.74
Constant		51.29		
\mathbb{R}^2		0.18		

Table 5: Logistic Regression of High Cardiovascular Fitness on Selected Independent Variables

	Odds Ratio	95% Confidence Interval	P value
Female	0.96	0.56-1.66	0.89
Age	1.05	1.02- 1.07	0.00
White	1.40	1.09- 1.77	0.01
Education	1.27	0.83-1.90	0.25
Body Mass Index	0.96	0.94-0.99	0.02
Serum Triglycerides	0.99	0.99-1.00	0.28
Serum Cotinine	1.00	0.99-1.00	0.69
Physical Activity	0.90	0.61-1.33	0.57
X^{2} (df)		18.08 (8)	

Figure 1. Conceptual Model for Determinants of Lung Function

