

Summer 8-9-2013

A Type-II Positive Allosteric Modulator of $\alpha 7$ nAChRs Reduces Brain Injury and Improves Neurological Function after Focal Cerebral Ischemia in Rats.

Fen Sun

University of North Texas Health Science Center at Fort Worth, Fen.Sun@unthsc.edu

Kunlin Jin

University of North Texas Health Science Center at Fort Worth, Kunlin.Jin@unthsc.edu

Victor Uteshev

University of North Texas Health Science Center at Fort Worth, Victor.Uteshev@unthsc.edu

Follow this and additional works at: http://digitalcommons.hsc.unt.edu/gsbs_facpubs

 Part of the [Medicine and Health Sciences Commons](#), and the [Neuroscience and Neurobiology Commons](#)

Recommended Citation

Sun, Fen; Jin, Kunlin; and Uteshev, Victor, "A Type-II Positive Allosteric Modulator of $\alpha 7$ nAChRs Reduces Brain Injury and Improves Neurological Function after Focal Cerebral Ischemia in Rats." (2013). *GSBS Faculty Scholarship*. Paper 5.
http://digitalcommons.hsc.unt.edu/gsbs_facpubs/5

This Article is brought to you for free and open access by the Graduate School of Biomedical Sciences at UNTHSC Scholarly Repository. It has been accepted for inclusion in GSBS Faculty Scholarship by an authorized administrator of UNTHSC Scholarly Repository. For more information, please contact Tom.Lyons@unthsc.edu.

A Type-II Positive Allosteric Modulator of $\alpha 7$ nAChRs Reduces Brain Injury and Improves Neurological Function after Focal Cerebral Ischemia in Rats

Fen Sun, Kunlin Jin, Victor V. Uteshev*

University of North Texas Health Science Center, Department of Pharmacology and Neuroscience, Fort Worth, TX, United States of America

Abstract

In the absence of clinically-efficacious therapies for ischemic stroke there is a critical need for development of new therapeutic concepts and approaches for prevention of brain injury secondary to cerebral ischemia. This study tests the hypothesis that administration of PNU-120596, a type-II positive allosteric modulator (PAM-II) of $\alpha 7$ nicotinic acetylcholine receptors (nAChRs), as long as 6 hours after the onset of focal cerebral ischemia significantly reduces brain injury and neurological deficits in an animal model of ischemic stroke. Focal cerebral ischemia was induced by a transient (90 min) middle cerebral artery occlusion (MCAO). Animals were then subdivided into two groups and injected intravenously (i.v.) 6 hours post-MCAO with either 1 mg/kg PNU-120596 (treated group) or vehicle only (untreated group). Measurements of cerebral infarct volumes and neurological behavioral tests were performed 24 hrs post-MCAO. PNU-120596 significantly reduced cerebral infarct volume and improved neurological function as evidenced by the results of Bederson, rolling cylinder and ladder rung walking tests. These results forecast a high therapeutic potential for PAMs-II as effective recruiters and activators of endogenous $\alpha 7$ nAChR-dependent cholinergic pathways to reduce brain injury and improve neurological function after cerebral ischemic stroke.

Citation: Sun F, Jin K, Uteshev VV (2013) A Type-II Positive Allosteric Modulator of $\alpha 7$ nAChRs Reduces Brain Injury and Improves Neurological Function after Focal Cerebral Ischemia in Rats. PLoS ONE 8(8): e73581. doi:10.1371/journal.pone.0073581

Editor: Alexander A. Mongin, Albany Medical College, United States of America

Received: June 18, 2013; **Accepted:** July 26, 2013; **Published:** August 9, 2013

Copyright: © 2013 Sun et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This study was supported by the National Institutes of Health Grant DK-082625 and the Start-Up funds from UNTHSC to V.V. Uteshev. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

* E-mail: Victor.Uteshev@unthsc.edu

Introduction

Clinical management of neuronal damage resulting from ischemic stroke generally involves only palliative treatments. Currently, the only FDA-approved drug therapy for ischemic stroke involves the intravenous use of tissue plasminogen activator (tPA) to dissolve clots [1]. This strategy appears to be effective in ischemic stroke, but only within the first 3 hours after the onset of ischemic stroke [2,3]. This strict limitation reduces the percent of stroke patients eligible for tPA to as low as ~2% [4]. Although in the last two decades substantial efforts have been invested in developing anti-ischemic medicine, these efforts have not resulted in clinically-efficacious therapies for ischemic stroke [5]. These failures highlight the need for development of new therapeutic concepts and approaches for prevention of brain injury secondary to ischemia. Among possible strategies, effective post-stroke treatments with broad therapeutic windows are likely to be the most valuable because of the unexpected nature of stroke. In this search, treatments that are based on recruiting and activating endogenous pathways receive special attention as these approaches are

expected to be highly efficacious and cause fewer adverse effects than approaches that utilize exogenous agents [6–8]. To complement these needs, this study evaluates neurological benefits of enhanced activation of $\alpha 7$ nicotinic acetylcholine receptors (nAChRs) by endogenous nicotinic agonists 6 hours after ischemic insult induced by middle cerebral artery occlusion (MCAO) in young adult rats.

There is a substantial body of supportive evidence linking age-, disease- and trauma-related reduction in the expression and function of $\alpha 7$ nAChRs to neurodegenerative, sensorimotor and psychiatric disorders associated with cognitive decline and attention deficits [9–24]. By contrast, activation of $\alpha 7$ nAChRs has been demonstrated to enhance neuronal resistance to ischemia and other insults in *in vivo*, *ex vivo* and *in vitro* experimental models [6,25–39], as well as improved cognitive performance in patients and animal models of neurodegenerative conditions including dementia, schizophrenia, brain trauma and aging [14,26,31,39–61]. An important rationale for the therapeutic use of $\alpha 7$ nAChR agents arises from the fact that $\alpha 7$ nAChRs are ubiquitously expressed throughout the brain [62] including brain regions that are highly

vulnerable to ischemia, such as cortex, striatum and hippocampus [63–66]. However, endogenous $\alpha 7$ nAChR agonists (i.e., choline and ACh) have not been regarded as potent therapeutic agents because physiological levels of choline/ACh do not appear to produce therapeutic levels of $\alpha 7$ activation [6]. This limitation has been recently resolved by the use of Type-II positive allosteric modulators (PAMs-II) of $\alpha 7$ nAChRs [6,8,48,67–73]. PAMs-II do not activate $\alpha 7$ nAChRs, but they inhibit desensitization and enhance $\alpha 7$ activation by nicotinic agonists, including endogenous choline and ACh [48,67,68]. Thus, PAMs-II only amplify activation of $\alpha 7$ nAChRs by endogenous nicotinic agonists released naturally as needed [8]. Accordingly, we have recently introduced a novel therapeutic paradigm [6] that converts endogenous choline/ACh into potent therapeutic agents for cerebral ischemia by enhancing activation of $\alpha 7$ nAChRs using PNU-120596, a PAM-II. In our previous proof-of-concept study [6], we have reported that a 3 hour pre-treatment with choline +PNU-120596 significantly delayed anoxic depolarization/injury of hippocampal CA1 pyramidal neurons in the complete oxygen/glucose deprivation model of ischemic stroke in acute hippocampal slices and activation of $\alpha 7$ nAChRs was required; while intravenous administration of PNU-120596 30 min post-ischemia in the MCAO model of ischemic stroke significantly reduced cerebral infarct volume [6]. The present study extends our previous findings and the therapeutic promise of PAMs-II by revealing that PNU-120596 reduces both the focal ischemia-induced cerebral infarct volume and neurological deficits even when administered as long as 6 hours after the ischemic onset. The results of this study further support the potential therapeutic utility of PAMs-II as effective recruiters and activators of endogenous $\alpha 7$ -dependent cholinergic pathways to reduce brain injury and improve neurological function secondary to focal cerebral ischemia.

Materials and Methods

Ethics Statement

Young adult male Sprague-Dawley (S.-D.) rats (~280 g) were used in experiments. The animal use was in accordance with the Guide for the Care and Use of Laboratory Animals (NIH 865-23, Bethesda, MD), and all experimental protocols were approved by the Institutional Animal Care and Use Committee of University of North Texas Health Science Center at Fort Worth, TX.

Animals

In total 22 animals were used in this study. Animals were housed 2 per tub in a Tecniplast Green Line IVC Sealsafe PLUS Rat rack on 1/8" corn cob bedding, with Envirodri shredded paper for enrichment. Animals were fed Purina Lab Diet 5LL2, and received filtered water via water bottles. Room lighting was kept below 50 Foot Candles (range of 30-40), and with a timer controlled 12:12 light dark cycle. Room temperature was maintained between 68–72 degrees, with humidity range of 30-70%. Cages were cleaned or changed at least once per week. The housing room contained only rats. The UNTHSC animal facility is AAALAC accredited and follows

or exceeds all of the requirements of the Guide for the Care and Use of Laboratory Animals.

Middle cerebral artery occlusion (MCAO)

Transient (90 min) focal cerebral ischemia was induced using the suture occlusion technique as previously described [74]. Animals (n=22; Charles River, Wilmington, MA) were anesthetized with 4% isoflurane mixed with 67% N₂O and 29% O₂ and delivered by a mask. After a midline incision in the neck, the left external carotid artery (ECA) was carefully exposed and dissected. A 19-mm, 4-0 monofilament nylon suture was inserted from the ECA into the left internal carotid artery to occlude the origin of left middle cerebral artery. After 90 min of occlusion, the thread was removed to allow reperfusion. The ECA was ligated, and the wound was closed. Rectal temperature was maintained at ~37° C using a heating pad.

A total of 22 animals were used in this study of which 1 animal from the control group died during the first hours of post-MCAO recovery prior to vehicle injections and another animal from the same control group died after vehicle injection, but prior to behavioral tests. Thus, the mortality rate was ~16.7% in the control group and 0% in the treatment group.

Drugs

PNU-120596 was obtained from the National Institute of Drug Addiction through the Research Resources Drug Supply Program as well as purchased from Selleck Chemicals (Houston, TX). Other chemicals were purchased from Sigma-Aldrich (St. Louis, MO).

PNU-120596

In all experiments of this study, 1 mg/kg PNU-120596 was administered via intravenous (i.v.) injections. Similar doses have been used in other studies [6,48,71,73]. To make a 50 mM stock solution (maximal achievable concentration is ~200 mM), PNU-120596 was dissolved in dimethyl sulfoxide (DMSO). The appropriate amounts of the stock solution (i.e., PNU+DMSO) or DMSO alone (i.e., vehicle) were injected as a single bolus. The amount of DMSO injected in each animal did not exceed 0.5 ml/kg.

Infarct Volume Measurements

Rats (n=10 per group) were anesthetized and euthanized by decapitation 24 hrs after MCAO. Brains were removed and coronal sections (2 mm thickness) immersed in 2% 2,3,5-triphenyltetrazolium chloride (TTC) in saline for 20 min at 37° C, then fixed for 2 hrs in 4% paraformaldehyde [75]. Infarct area, left hemisphere area, and total brain area were measured by a blinded observer using the ImageJ software, and areas were multiplied by the distance between sections to obtain the respective volumes. Infarct volume was calculated as a percentage of the volume of the contralateral hemisphere, as described previously [76].

Neurobehavioral testing

Rats (n=10 per group) underwent neurobehavioral tests to evaluate functional outcome of treatments with PNU-120596. Animals were trained prior to MCAO (training period: 3 days, 3 trials per day) and deficits were assessed 24 hrs thereafter. The order of testing (Bederson→cylinder→ladder rung walking) was always the same to keep the testing conditions identical for all animals. Although it is unlikely that subjecting animals to early tests in the sequence facilitated or inhibited the animal performance in the later tests, we cannot completely rule out a possibility of inter-test interactions.

Bederson test

Bederson score was used to assess the neurological deficit using a four-level scale [77]: 0, normal; 1, forelimb flexion; 2, decreased resistance to lateral push; 3, circling.

Cylinder Test

Forelimb use bias was analyzed by observing the rat's movements over 3-minute intervals in a transparent, 18-cm-wide, 30-cm-high poly(methyl methacrylate) cylinder. A mirror behind the cylinder made it possible to observe and record forelimb movements when the rat was facing away from the examiner. After an episode of rearing and wall exploration, a landing was scored for the first limb to contact the wall or for both limbs if they made simultaneous contact. Percentage use of the impaired limb was calculated.

Ladder rung walking test

The ladder rung walking test is sensitive for quantifying skilled locomotion. The degree of motor dysfunction after MCAO was measured by counting the number of foot-faults of the impaired limbs per round, as described previously [78]. Baseline and post-operative testing sessions consisted of three traverses across the ladder. An error was scored for any foot slip or misstep. The number of errors of the affected forelimb and hindlimb in each trial was counted. The mean number of errors in three traverses was calculated.

Statistical Analysis

Statistical significance of differences among groups was defined by the p-value (i.e., * p<0.05; *** p<0.001) using the two-tailed Mann–Whitney U-test. A non-parametric Mann–Whitney U-test was used because this study did not assume any specific underlying distribution (i.e., Gaussian) of data and had a relatively small sample size (n=5-10). We recognize that non-parametric statistics are often less powerful than parametric statistics and thus, more prone to Type-II error (i.e., missing significance when it is present) [79]. However, in this particular study, differences among groups have been found significant in all experiments further supporting our conclusions. The results are presented as mean±S.E.M.

Results

PNU-120596 significantly reduces cerebral infarct volume

In the group of animals defined as treated, PNU-120596 (1 mg/kg) was administered intravenously (i.v.) 6 hrs post-MCAO and the effects of PNU-120596 on cerebral infarct volume were evaluated 24 hrs post-MCAO using the TTC staining (see Methods). In the matching control group of animals only vehicle (i.e., DMSO) was administered via i.v. injections. Only the left MCA was occluded in each experiment. The results of these experiments demonstrated significant reduction in the infarct volume of treated vs. untreated animals (two-tailed, Mann–Whitney U-test): p=0.0147 (n=10; Figure 1).

PNU-120596 significantly improves neurological performance post-MCAO

The same treated and untreated animals that were used for histological measurements (Figure 1) were used in behavioral experiments 15 min prior to the animal anesthesia/euthanasia and brain tissue collection for histology (i.e., ~24 hrs post-MCAO). PNU-120596 significantly improved neurological function of treated (n=10) vs. untreated (n=10) animals as evidenced by the results of the following behavioral tests (two-tailed, Mann–Whitney U-test): Bederson (p=0.0385; Figure 2A), rolling cylinder (p=0.0124; Figure 2B), ladder rung walking (forelimb) (p=0.0486; Figure 2C) and ladder rung walking (hindlimb) (p=0.0007; Figure 2D). Therefore, the results of these experiments convincingly demonstrate that PNU-120596 produces significant neurological benefits even when it is administered as long as 6 hrs post-MCAO.

Discussion

The key finding of this study is that PNU-120596, a previously reported highly selective PAM-II of $\alpha 7$ nAChRs, significantly reduces cerebral infarct volume and neurological deficits in the MCAO model of ischemic stroke in rats when the drug is administered as long as 6 hrs post-MCAO. Such a remarkable persistent post-MCAO effectiveness of PNU-120596 invites more comprehensive pre-clinical studies of the PAM-II class of compounds aiming at giving health care providers an effective tool to reducing brain injury and improving neurological function secondary to cerebral ischemic stroke hours after the initial ischemic event. The therapeutic benefits produced by PNU-120596 originate from its ability to convert endogenous agonists of $\alpha 7$ nAChRs (i.e., choline and ACh) into highly potent therapeutic agents [6,48,67,68]. Thus, PAMs-II may create a conceptually novel family of treatments that are based on a novel and substantively different mechanism, i.e., recruiting and activating endogenous $\alpha 7$ -dependent cholinergic pathways. Treatments that incorporate endogenous compounds and mechanisms are expected to be highly efficacious and cause fewer adverse effects than treatments that utilize exogenous agents.

These results extend our previous findings that demonstrated a high therapeutic efficacy of PNU-120596 administered intravenously 30 min after focal cerebral ischemia




Figure 1. PNU-120596 significantly reduces the size of brain injury induced by focal cerebral ischemia. Focal cerebral ischemia was induced by a transient (90 min) middle cerebral artery occlusion (MCAO). Then, 6 hrs post-MCAO, animals were given i.v. injections of either 1 mg/kg PNU-120596 dissolved in DMSO at 50 mM (i.e., treated group; n=10) or the matched amount of DMSO only (i.e., untreated group; n=10). Typical examples of injured whole-brain coronal sections (2 mm thick) obtained from untreated (i.e., DMSO only) (**A**) and treated (1 mg/kg PNU-120596) (**B**) animals. Treated and untreated animals were anesthetized and euthanized 24 hrs after MCAO (i.e., 18 hrs after PNU-120596 or DMSO injections) and brain sections were prepared for histological analysis. **C**) A summary: MCAO-induced infarct volumes were significantly smaller in treated vs. untreated animals: $p=0.0147$ (n=10; two-tailed, the Mann–Whitney U-test). The results are presented as mean \pm S.E.M.

doi: 10.1371/journal.pone.0073581.g001




Figure 2. PNU-120596 significantly improves neurological function after focal cerebral ischemia. The same treated (n=10) and untreated (n=10) animals that were used for histological analysis (Figure 1) were subjected to neurological tests 15 min prior to anesthesia/euthanasia and collection of brain sections for histological analysis. PNU-120596 significantly improved neurological function post-MCAO in treated (n=10) vs. untreated (n=10) groups of animals as evidenced by the results of the following tests (two-tailed, the Mann–Whitney U-test): **A)** Bederson, (p=0.0385); **B)** Rolling cylinder, (p=0.0124); **C)** Ladder rung walk, (forelimb), (p=0.0486); and **D)** Ladder rung walk, (hindlimb), (p=0.0007). The results are presented as mean±S.E.M.

doi: 10.1371/journal.pone.0073581.g002

[6]. Intriguingly, infarct volumes measured in animals treated with PNU-120596 30 min (n=5 [6]) and 6 hrs (n=10; this study) post-MCAO were not significantly different (p=0.2404, two-tailed Mann-Whitney test). Similarly, the corresponding infarct volumes measured in untreated animals (i.e., DMSO only) 30 min (n=5 [6]) and 6 hrs (n=10; this study) post-MCAO were also not significantly different (p=0.5921, two-tailed Mann-Whitney test). Therefore, it is likely that the therapeutic efficacy of PNU-120596 extends considerably beyond the 6 hrs post-ischemic delay tested in this study. By contrast, the therapeutic efficacy of donepezil, an inhibitor of ACh hydrolysis, has been reported to cease within the first 2 hrs post-MCAO [80]. Although the reason for these differences is not known, it may be related to the ability of PNU-120596 to inhibit $\alpha 7$ desensitization and thus, generate persistent $\alpha 7$ nAChR-mediated currents in the presence of physiological/endogenous choline [67–69] even though these currents appear to be reduced at physiological temperatures [81]. By inhibiting ACh hydrolysis, donepezil elevates the extracellular levels of ACh (a non-selective agonist of nicotinic and muscarinic AChRs), but does not appear to produce therapeutic levels of nicotinic and muscarinic AChR activation after 1–2 hrs post-ischemia [80].

The therapeutic utility of PAM-II-based strategies is supported by the ubiquitous expression of $\alpha 7$ nAChRs in the brain and especially, in the brain regions that are selectively vulnerable to ischemia, such as cortex, striatum and hippocampus [63–66]. Activation of $\alpha 7$ nAChRs has been shown to enhance neuronal resistance to ischemic and other types of insults [6,31,32,38,39,63,82] as well as improve cognitive performance in patients and animal models of schizophrenia [49,72,73,83], dementia [56,61,84] and traumatic brain injury [39]. Moreover, PNU-120596 has been recently demonstrated to produce robust anti-nociceptive effects by enhancing the potency of endogenous choline for $\alpha 7$ nAChR activation [70,71]. Although choline is a full selective endogenous $\alpha 7$ nAChR agonist, near-physiological levels of choline (i.e., ~20 μ M) [12,85–87] are sub-threshold for $\alpha 7$ activation (EC_{50} ~0.5 mM) [88] and tend to induce $\alpha 7$ desensitization (IC_{50} ~40 μ M) [87]. These limitations can be overcome by the use of PAMs-II, such as PNU-120596. PNU-120596 inhibits $\alpha 7$ desensitization and increases the potency of endogenous choline/ACh for $\alpha 7$ activation producing a weak persistent and tunable activation of $\alpha 7$ nAChRs [67–69] – an activation modality of $\alpha 7$ nAChRs that can benefit neuronal survival as discussed previously [6,27,31,32,38,39,63,82]. Moreover, energy deprivation and cell death/dysfunction can considerably elevate the concentration of choline in the extracellular space [89–91] providing a large source of this endogenous $\alpha 7$ nAChR agonist as has been recently demonstrated by direct measurements of choline/ACh levels in the ischemic core and penumbra in the MCAO model of ischemic stroke in rats [92]. It is intriguing to hypothesize that these elevated levels of choline near the site of injury may robustly enhance neuronal resistance to ischemic injury, while PNU-120596 augments this endogenous therapeutic process [6].

Although the exact cellular and molecular mechanisms of the therapeutic effects of PNU-120596 are not known, $\alpha 7$ nAChR-

mediated Ca^{2+} -dependent activation of JAK2/AKT-dependent pathways are likely candidates [82,93–95]. These likely mechanisms would be expected to delay mitochondrial dysfunction and thus, PNU-120596-treated neurons may be able to better meet the energy demand of ischemic/hypoglycemic conditions and significantly delay the ultimate failure of the Na^+/K^+ -ATPase pumps. Such a failure would cause a rapid loss of the neuronal trans-membrane electrochemical gradient leading to transient or terminal anoxic depolarization [6]. It may seem counterintuitive that excitatory currents (i.e., $\alpha 7$ nAChR-mediated) could delay anoxic depolarization and reduce brain injury [6]. However, this concept reflects a common motif in how central neurons respond to insults, i.e., the existence of optimal neuroprotective levels and spatiotemporal patterns of cytosolic Ca^{2+} elevations [8,14,27,32,96–101]. While sub-optimal levels of cytosolic Ca^{2+} are ineffective, excessive Ca^{2+} influx is toxic. By contrast, moderate elevations in cytosolic Ca^{2+} levels, for example, via a K^+ -induced depolarization or weak persistent activation of highly Ca^{2+} -permeable $\alpha 7$ nAChRs [102–104] have been shown to protect neurons from injury in a variety of toxicity/insult models [6,27,28,32,33,38,98,105,106]. These therapeutic levels of $\alpha 7$ nAChR activation are consistent with the weak persistent modality of $\alpha 7$ nAChR activation generated by physiological concentrations of choline in the presence of PNU-120596 [67–69].

Moreover, the reported therapeutic efficacy of PNU-120596 may have resulted, at least in part, from enhanced activation of $\alpha 7$ nAChRs expressed in the autonomic neuronal circuitry which may have provided a neurogenic (e.g., adrenergic, nitrenergic [107,108]) control over vascular tone and collateral blood circulation. In addition, functional $\alpha 7$ nAChRs are expressed in numerous non-neuronal tissues including glial [109–111] and immune cells [112–114]. Thus, several therapeutic components of $\alpha 7$ nAChR activation in multiple neuronal and non-neuronal tissues may have contributed to the significant therapeutic efficacy of PNU-120596 reported in this and previous *in vivo* studies [6,70–73,114,115]. These potential individual sources of brain protection and their relative contributions to the therapeutic effects of PNU-120596 are not known and present great interest.

One potential limitation of this study is that it does not include experiments with $\alpha 7$ nAChR antagonists (e.g., methyllycaconitine; MLA). Although PNU-120596 is highly selective for $\alpha 7$ nAChRs and to-date non- $\alpha 7$ -mediated effects of PNU-120596 have not been reported, there is a slight chance that PNU-120596 activates both $\alpha 7$ -dependent and yet unknown, $\alpha 7$ -independent pathways. In that unlikely event, the use of highly selective $\alpha 7$ nAChR antagonists would be critical for distinguishing among $\alpha 7$ -dependent and $\alpha 7$ -independent components of the effects of PNU-120596. However, experiments using MLA *in vivo* may not be straightforward as evidenced from a previous report where the effects of MLA on certain behavioral functions were bell-shaped [116]. Thus, a

series of positive and negative controls will need to be conducted using selective $\alpha 7$ agonists (e.g., DMXBA; 3-(2,4-dimethoxybenzylidene)-anabaseine, also known as GTS-21) to determine the effective regimens of MLA as applicable to MCAO. This work has not yet been done in this laboratory.

Another possible limitation is that we have not tested the effects of PNU-120596 on neurological performance of control (sham) animals (i.e., in the absence of MCAO-induced injury). This is because control animals perform these tests nearly flawlessly leaving no room for significant improvement by PNU-120596. However, because of this limitation we cannot exclude the possibility that PNU-120596 is a performance enhancing drug which is also effective in the absence of MCAO-induced injury and thus, the therapeutic efficacy of PNU-120596 post-MCAO may not be directly related to MCAO-induced injury, but extends the performance-enhancing potential of PNU-120596 in the absence of injury.

Certain genetic, age- and trauma-related neurodegenerative, sensorimotor, and psychiatric disorders characterized by cognitive decline and attention deficits (e.g., schizophrenia, dementia and traumatic brain injury) are directly associated with decreased cholinergic tone and a decrease, but not disappearance, of functional $\alpha 7$ nAChRs [10,49,117]. By increasing and partially restoring $\alpha 7$ -dependent cholinergic tone, PAMs-II would be expected to improve cognitive function and attention impairments in these patients and animal models [39,49,53,56,61,84]. In this regard, treatments with PNU-120596 or functionally-similar PAMs-II compounds may benefit individuals with ischemic stroke and certain age- and trauma-related cognitive deficits via multiple mechanisms and routes of action.

In conclusion, this study demonstrates a remarkable reduction in the size of cerebral injury and significant improvements in neurological function upon intravenous administration of PNU-120596 as long as 6 hours after the onset of transient focal cerebral ischemia. These results further support the potential therapeutic utility of PAMs-II as effective recruiters and activators of endogenous $\alpha 7$ -dependent cholinergic pathways and extend the therapeutic promise of this novel class of compounds.

Acknowledgements

We thank the National Institute on Drug Abuse Research Resources Drug Supply Program for PNU-120596.

Author Contributions

Conceived and designed the experiments: FS KJ VU. Performed the experiments: FS. Analyzed the data: FS KJ VU. Contributed reagents/materials/analysis tools: KJ VU. Wrote the manuscript: VU. Interpreted results: FS KJ VU. Edited and revised manuscript: KJ VU. Prepared figures: FS VU.

References

- Furlan AJ (2002) Acute stroke therapy: beyond i.v. tPA. *Cleve Clin J Med* 69: 730-734. doi:10.3949/ccjm.69.9.730. PubMed: 12222978.
- Adams HP Jr., del Zoppo G, Alberts MJ, Bhatt DL, Brass L et al. (2007) Guidelines for the early management of adults with ischemic stroke: a guideline from the American Heart Association/American Stroke Association Stroke Council, Clinical Cardiology Council, Cardiovascular Radiology and Intervention Council, and the Atherosclerotic Peripheral Vascular Disease and Quality of Care Outcomes in Research Interdisciplinary Working Groups: the American Academy of Neurology affirms the value of this guideline as an educational tool for neurologists. *Stroke* 38: 1655-1711. doi:10.1161/STROKEAHA.107.181486. PubMed: 17431204.
- Brott T, Bogousslavsky J (2000) Treatment of acute ischemic stroke. *N Engl J Med* 343: 710-722. doi:10.1056/NEJM200009073431007. PubMed: 10974136.
- Katzan IL, Furlan AJ, Lloyd LE, Frank JL, Harper DL et al. (2000) Use of tissue-type plasminogen activator for acute ischemic stroke: the Cleveland area experience. *JAMA* 283: 1151-1158. doi:10.1001/jama.283.9.1151. PubMed: 10703777.
- Richard Green A, Odergren T, Ashwood T (2003) Animal models of stroke: do they have value for discovering neuroprotective agents? *Trends Pharmacol Sci* 24: 402-408. doi:10.1016/S0165-6147(03)00192-5. PubMed: 12915049.
- Kalappa BI, Sun F, Johnson SR, Jin K, Uteshev VV (2013) A positive allosteric modulator of alpha7 nAChRs augments neuroprotective effects of endogenous nicotinic agonists in cerebral ischemia. *Br J Pharmacol*.
- Lo EH (2008) A new penumbra: transitioning from injury into repair after stroke. *Nat Med* 14: 497-500. doi:10.1038/nm1735. PubMed: 18463660.
- Uteshev VV (2012) Alpha 7 Nicotinic ACh Receptors as a Ligand-Gated Source of Ca(2+) Ions: The Search for a Ca (2+) Optimum. *Adv Exp Med Biol* 740: 603-638.
- Jenden DJ, Scremin OU, Roch M, Li G (1996) The influence of aging on whole body choline release and clearance. *Life Sci* 58: 2003-2009. doi:10.1016/0024-3205(96)00191-9. PubMed: 8637430.
- Guan ZZ, Zhang X, Ravid R, Nordberg A (2000) Decreased protein levels of nicotinic receptor subunits in the hippocampus and temporal cortex of patients with Alzheimer's disease. *JNeurochem* 74: 237-243.
- Nordberg A (2001) Nicotinic receptor abnormalities of Alzheimer's disease: therapeutic implications. *Biol Psychiatry* 49: 200-210. doi:10.1016/S0006-3223(00)01125-2. PubMed: 11230871.
- Sarter M, Parikh V (2005) Choline transporters, cholinergic transmission and cognition. *Nat Rev Neurosci* 6: 48-56. doi:10.1038/nrn1588. PubMed: 15611726.
- Martin LF, Freedman R (2007) Schizophrenia and the alpha7 Nicotinic Acetylcholine Receptor. *Int Rev Neurobiol* 78: 225-246. doi:10.1016/S0074-7742(06)78008-4. PubMed: 17349863.
- Thomsen MS, Hansen HH, Timmerman DB, Mikkelsen JD (2010) Cognitive improvement by activation of alpha7 nicotinic acetylcholine receptors: from animal models to human pathophysiology. *Curr Pharm Des* 16: 323-343. doi:10.2174/138161210790170094. PubMed: 20109142.
- Leonard S, Breese C, Adams C, Benhammou K, Gault J et al. (2000) Smoking and schizophrenia: abnormal nicotinic receptor expression. *Eur J Pharmacol* 393: 237-242. PubMed: 10771019.
- Freedman R, Adams CE, Leonard S (2000) The alpha7-nicotinic acetylcholine receptor and the pathology of hippocampal interneurons in schizophrenia. *J Chem Neuroanat* 20: 299-306. doi:10.1016/S0891-0618(00)00109-5. PubMed: 11207427.
- Stevens KE, Freedman R, Collins AC, Hall M, Leonard S et al. (1996) Genetic correlation of inhibitory gating of hippocampal auditory evoked response and alpha-bungarotoxin-binding nicotinic cholinergic receptors in inbred mouse strains. *Neuropsychopharmacology* 15: 152-162. doi:10.1016/0893-133X(95)00178-G. PubMed: 8840351.
- Felix R, Levin ED (1997) Nicotinic antagonist administration into the ventral hippocampus and spatial working memory in rats. *Neuroscience* 81: 1009-1017. doi:10.1016/S0306-4522(97)00224-8. PubMed: 9330363.
- Wevers A, Witter B, Moser N, Burghaus L, Banerjee C et al. (2000) Classical Alzheimer features and cholinergic dysfunction: towards a unifying hypothesis? *Acta Neurol Scand Suppl* 176: 42-48. PubMed: 11261804.
- Freedman R, Hall M, Adler LE, Leonard S (1995) Evidence in postmortem brain tissue for decreased numbers of hippocampal nicotinic receptors in schizophrenia. *Biol Psychiatry* 38: 22-33. doi:10.1016/0006-3223(94)00252-X. PubMed: 7548469.
- Perry EK, Morris CM, Court JA, Cheng A, Fairbairn AF et al. (1995) Alteration in nicotine binding sites in Parkinson's disease, Lewy body dementia and Alzheimer's disease: possible index of early neuropathology. *Neuroscience* 64: 385-395. doi:10.1016/0306-4522(94)00410-7. PubMed: 7700528.
- Nordberg A, Winblad B (1986) Reduced number of [3H]nicotine and [3H]acetylcholine binding sites in the frontal cortex of Alzheimer brains. *Neurosci Lett* 72: 115-119. doi:10.1016/0304-3940(86)90629-4. PubMed: 3808458.
- Shimohama S, Taniguchi T, Fujiwara M, Kameyama M (1986) Changes in nicotinic and muscarinic cholinergic receptors in Alzheimer-type dementia. *J Neurochem* 46: 288-293. doi:10.1111/j.1471-4159.1986.tb12960.x. PubMed: 3940287.
- London ED, Ball MJ, Waller SB (1989) Nicotinic binding sites in cerebral cortex and hippocampus in Alzheimer's dementia. *Neurochem Res* 14: 745-750. doi:10.1007/BF00964952. PubMed: 2812250.
- Ren K, King MA, Liu J, Siemann J, Altman M et al. (2007) Alpha 7 nicotinic receptor agonist 4OH-GTS-21 protects axotomized septohippocampal cholinergic neurons in wild type but not amyloid-overexpressing transgenic mice. *Neuroscience* 148: 230-237.
- Takeuchi H, Yanagida T, Inden M, Takata K, Kitamura Y, et al. (2009) Nicotinic receptor stimulation protects nigral dopaminergic neurons in rotenone-induced Parkinson's disease models. *J Neurosci Res* 87: 576-585.
- Shimohama S, Greenwald DL, Shafron DH, Akaika A, Maeda T et al. (1998) Nicotinic alpha 7 receptors protect against glutamate neurotoxicity and neuronal ischemic damage. *Brain Res* 779: 359-363. doi:10.1016/S0006-8993(97)00194-7. PubMed: 9473725.
- Akaika A, Tamura Y, Yokota T, Shimohama S, Kimura J (1994) Nicotine-induced protection of cultured cortical neurons against N-methyl-D-aspartate receptor-mediated glutamate cytotoxicity. *Brain Res* 644: 181-187. doi:10.1016/0006-8993(94)91678-0. PubMed: 7519524.
- Kaneko S, Maeda T, Kume T, Kochiyama H, Akaika A et al. (1997) Nicotine protects cultured cortical neurons against glutamate-induced cytotoxicity via alpha7-neuronal receptors and neuronal CNS receptors. *Brain Res* 765: 135-140. doi:10.1016/S0006-8993(97)00556-8. PubMed: 9310404.
- Kihara T, Shimohama S, Sawada H, Kimura J, Kume T et al. (1997) Nicotinic receptor stimulation protects neurons against beta-amyloid toxicity. *Ann Neurol* 42: 159-163. doi:10.1002/ana.410420205. PubMed: 9266724.
- Meyer EM, Tay ET, Zoltewicz JA, Papke RL, Meyers C et al. (1998) Neuroprotective and memory-related actions of novel $\alpha 7$ nicotinic agents with different mixed agonist/antagonist properties. *J PharmacolExpTher* 284: 1026-1032.
- Li Y, Papke RL, He YJ, Millard WJ, Meyer EM (1999) Characterization of the neuroprotective and toxic effects of alpha7 nicotinic receptor activation in PC12 cells. *Brain Res* 830: 218-225. doi:10.1016/S0006-8993(99)01372-4. PubMed: 10366678.
- Shimohama S, Kihara T (2001) Nicotinic receptor-mediated protection against beta-amyloid neurotoxicity. *Biol Psychiatry* 49: 233-239. doi:10.1016/S0006-3223(00)01100-8. PubMed: 11230874.
- Verbois SL, Scheff SW, Pauly JR (2003) Chronic nicotine treatment attenuates alpha 7 nicotinic receptor deficits following traumatic brain injury. *Neuropharmacology* 44: 224-233. doi:10.1016/S0028-3908(02)00366-0. PubMed: 12623221.
- Buccafusco JJ (2004) Neuronal Nicotinic Receptor Subtypes: DEFINING THERAPEUTIC TARGETS. *Mol Interv* 4: 285-295. doi:10.1124/mi.4.5.8. PubMed: 15471911.
- Fucile S, Renzi M, Lauro C, Limatola C, Ciotti T et al. (2004) Nicotinic cholinergic stimulation promotes survival and reduces motility of cultured rat cerebellar granule cells. *Neuroscience* 127: 53-61. doi:10.1016/j.neuroscience.2004.04.017. PubMed: 15219668.
- Rosa AO, Egea J, Gandia L, López MG, García AG (2006) Neuroprotection by nicotine in hippocampal slices subjected to oxygen-glucose deprivation: involvement of the alpha7 nAChR subtype. *J Mol Neurosci* 30: 61-62. doi:10.1385/JMN:30:1:61. PubMed: 17192628.
- Egea J, Rosa AO, Sobrado M, Gandia L, López MG et al. (2007) Neuroprotection afforded by nicotine against oxygen and glucose deprivation in hippocampal slices is lost in alpha7 nicotinic receptor knockout mice. *Neuroscience* 145: 866-872. doi:10.1016/j.neuroscience.2006.12.036. PubMed: 17291692.
- Guseva MV, Hopkins DM, Scheff SW, Pauly JR (2008) Dietary choline supplementation improves behavioral, histological, and neurochemical outcomes in a rat model of traumatic brain injury. *Neurotrauma* 25: 975-983. doi:10.1089/neu.2008.0516.

40. Buccafusco JJ, Letchworth SR, Bencherif M, Lippiello PM (2005) Long-lasting cognitive improvement with nicotinic receptor agonists: mechanisms of pharmacokinetic-pharmacodynamic discordance. *Trends Pharmacol Sci* 26: 352-360. doi:10.1016/j.tips.2005.05.007. PubMed: 15946748.
41. Buccafusco JJ, Terry AV Jr., Decker MW, Gopalakrishnan M (2007) Profile of nicotinic acetylcholine receptor agonists ABT-594 and A-582941, with differential subtype selectivity, on delayed matching accuracy by young monkeys. *Biochem Pharmacol* 74: 1202-1211. doi: 10.1016/j.bcp.2007.07.010. PubMed: 17706609.
42. Kitagawa H, Takenouchi T, Azuma R, Wesnes KA, Kramer WG et al. (2003) Safety, pharmacokinetics, and effects on cognitive function of multiple doses of GTS-21 in healthy, male volunteers. *Neuropsychopharmacology* 28: 542-551. doi:10.1038/sj.npp.1300028. PubMed: 12629535.
43. Leiser SC, Bowlby MR, Comery TA, Dunlop J (2009) A cog in cognition: how the alpha 7 nicotinic acetylcholine receptor is geared towards improving cognitive deficits. *Pharmacol Ther* 122: 302-311. doi:10.1016/j.pharmthera.2009.03.009. PubMed: 19351547.
44. Olincy A, Stevens KE (2007) Treating schizophrenia symptoms with an alpha7 nicotinic agonist, from mice to men. *Biochem Pharmacol* 74: 1192-1201. doi:10.1016/j.bcp.2007.07.015. PubMed: 17714692.
45. Arendash GW, Sengstock GJ, Sanberg PR, Kem WR (1995) Improved learning and memory in aged rats with chronic administration of the nicotinic receptor agonist GTS-21. *Brain Res* 674: 252-259. doi: 10.1016/0006-8993(94)01449-R. PubMed: 7796104.
46. Meyer EM, Tay ET, Papke RL, Meyers C, Huang GL et al. (1997) Effects of 3-[2,4-dimethoxybenzylidene]anabaseine (DMXB) on rat nicotinic receptors and memory-related behaviors. *Brain Res* 768: 49-56. doi:10.1016/S0006-8993(97)00536-2. PubMed: 9369300.
47. Ross RG, Stevens KE, Proctor WR, Leonard S, Kiskey MA et al. (2010) Research review: Cholinergic mechanisms, early brain development, and risk for schizophrenia. *J Child Psychol Psychiatry* 51: 535-549. doi: 10.1111/j.1469-7610.2009.02187.x. PubMed: 19925602.
48. Hurst RS, Hajos M, Raggenbass M, Wall TM, Higdon NR et al. (2005) A novel positive allosteric modulator of the alpha7 neuronal nicotinic acetylcholine receptor: in vitro and in vivo characterization. *J Neurosci* 25: 4396-4405.
49. Olincy A, Harris JG, Johnson LL, Pender V, Kongs S et al. (2006) Proof-of-concept trial of an alpha7 nicotinic agonist in schizophrenia. *Arch Gen Psychiatry* 63: 630-638. doi:10.1001/archpsyc.63.6.630. PubMed: 16754836.
50. Martin EJ, Panikar KS, King MA, Deyrup M, Hunter B et al. (1994) Cytoprotective actions of 2,4-dimethoxybenzylidene anabaseine in differentiated PC12 cells and septal cholinergic cells. *Drug Dev Res* 31: 134-141.
51. Briggs CA, Anderson DJ, Brioni JD, Buccafusco JJ, Buckley MJ et al. (1997) Functional characterization of the novel neuronal nicotinic acetylcholine receptor ligand GTS-21 in vitro and in vivo. *Pharmacol Biochem Behav* 57: 231-241. doi:10.1016/S0091-3057(96)00354-1. PubMed: 9164577.
52. Van Kampen M, Selbach K, Schneider R, Schiegel E, Boess F et al. (2004) AR-R 17779 improves social recognition in rats by activation of nicotinic alpha7 receptors. *Psychopharmacology (Berl)* 172: 375-383. doi:10.1007/s00213-003-1668-7. PubMed: 14727003.
53. Brown KL, Comalli DM, De Biasi M, Woodruff-Pak DS (2010) Trace eyeblink conditioning is impaired in $\alpha 7$ but not in $\beta 2$ nicotinic acetylcholine receptor knockout mice. *Front Behav Neurosci* 4: 166. PubMed: 20976039.
54. Woodruff-Pak DS, Li YT, Kem WR (1994) A nicotinic agonist (GTS-21), eyeblink classical conditioning, and nicotinic receptor binding in rabbit brain. *Brain Res* 645: 309-317. doi:10.1016/0006-8993(94)91665-9. PubMed: 8062092.
55. Wishka DG, Walker DP, Yates KM, Reitz SC, Jia S et al. (2006) Discovery of N-[(p. 3R)-1-azabicyclo[2.2.2]oct-3-yl]furo[2,3-c]pyridine-5-carboxamide, an agonist of the alpha7 nicotinic acetylcholine receptor, for the potential treatment of cognitive deficits in schizophrenia: synthesis and structure-activity relationship. *J Med Chem* 49: 4425-4436.
56. Bitner RS, Bunnelle WH, Decker MW, Drescher KU, Kohlhaas KL et al. (2010) In vivo pharmacological characterization of a novel selective alpha7 neuronal nicotinic acetylcholine receptor agonist ABT-107: preclinical considerations in Alzheimer's disease. *J Pharmacol Exp Ther* 334: 875-886. doi:10.1124/jpet.110.167213. PubMed: 20504913.
57. Bitner RS, Bunnelle WH, Anderson DJ, Briggs CA, Buccafusco J et al. (2007) Broad-spectrum efficacy across cognitive domains by alpha7 nicotinic acetylcholine receptor agonist correlates with activation of ERK1/2 and CREB phosphorylation pathways. *J Neurosci* 27: 10578-10587. doi:10.1523/JNEUROSCI.2444-07.2007. PubMed: 17898229.
58. Boess FG, De Vry J, Erb C, Flessner T, Hendrix M et al. (2007) The novel alpha7 nicotinic acetylcholine receptor partial agonist N-[(p. 3R)-1-azabicyclo[2.2.2]oct-3-yl]-7-[2-(methoxy)phenyl]-1-benzofuran-2-carboxamide improves working and recognition memory in rodents. *J Pharmacol Exp Ther* 321: 716-725.
59. Pichat P, Bergis OE, Terranova JP, Urani A, Duarte C et al. (2007) SSR180711, a novel selective alpha7 nicotinic receptor partial agonist: (II) efficacy in experimental models predictive of activity against cognitive symptoms of schizophrenia. *Neuropsychopharmacology* 32: 17-34. doi:10.1038/sj.npp.1301188. PubMed: 16936709.
60. Tatsumi R, Fujio M, Takanashi S, Numata A, Katayama J et al. (2006) (R)-3'-(3-methylbenzo[b]thiophen-5-yl)spiro[1-azabicyclo[2.2.2]octane-3,5'-oxazolidin]-2'-one, a novel and potent alpha7 nicotinic acetylcholine receptor partial agonist displays cognitive enhancing properties. *J Med Chem* 49: 4374-4383. doi:10.1021/jm060249c. PubMed: 16821797.
61. Ren K, Thinschmidt J, Liu J, Ai L, Papke RL et al. (2007) Alpha 7 Nicotinic receptor gene delivery into mouse hippocampal neurons leads to functional receptor expression, improved spatial memory-related performance, and tau hyperphosphorylation. *Neuroscience* 145: 314-322.
62. Clarke PBS, Schwartz RD, Paul SM, Pert CB, Pert A (1985) Nicotinic binding in rat brain: autoradiographic comparison of [3 H] acetylcholine [3 H] nicotine and [125 I]-alpha-bungarotoxin. *J Neurosci* 5: 1307-1315.
63. Quik M, Huang LZ, Parameswaran N, Bordia T, Campos C et al. (2009) Multiple roles for nicotine in Parkinson's disease. *Biochem Pharmacol* 78: 677-685. doi:10.1016/j.bcp.2009.05.003. PubMed: 19433069.
64. Breese CR, Adams C, Logel J, Drebing C, Rollins Y et al. (1997) Comparison of the regional expression of nicotinic acetylcholine receptor alpha7 mRNA and [125I]-alpha-bungarotoxin binding in human postmortem brain. *J Comp Neurol* 387: 385-398. doi:10.1002/(SICI)1096-9861(19971027)387:3. PubMed: 9335422.
65. Whiteaker P, Davies AR, Marks MJ, Blagbrough IS, Potter BV et al. (1999) An autoradiographic study of the distribution of binding sites for the novel alpha7-selective nicotinic radioligand [3H]-methyllycaconitine in the mouse brain. *Eur J Neurosci* 11: 2689-2696. doi:10.1046/j.1460-9568.1999.00685.x. PubMed: 10457165.
66. Woodruff TM, Thundiyil J, Tang SC, Sobey CG, Taylor SM et al. (2011) Pathophysiology, treatment, and animal and cellular models of human ischemic stroke. *Mol Neurodegener* 6: 11. doi:10.1186/1750-1326-6-11. PubMed: 21266064.
67. Gusev AG, Uteshev VV (2010) Physiological concentrations of choline activate native alpha7-containing nicotinic acetylcholine receptors in the presence of PNU-120596 [1-(5-chloro-2,4-dimethoxyphenyl)-3-(5-methylisoxazol-3-yl)-urea]. *J Pharmacol Exp Ther* 332: 588-598. doi: 10.1124/jpet.109.162099. PubMed: 19923442.
68. Kalappa BI, Gusev AG, Uteshev VV (2010) Activation of functional $\alpha 7$ -containing nAChRs in hippocampal CA1 pyramidal neurons by physiological levels of choline in the presence of PNU-120596. *PLOS ONE* 5: e13964. doi:10.1371/journal.pone.0013964. PubMed: 21103043.
69. Uteshev VV (2012) Somatic integration of single ion channel responses of $\alpha 7$ nicotinic acetylcholine receptors enhanced by PNU-120596. *PLOS ONE* 7: e32951. doi:10.1371/journal.pone.0032951. PubMed: 22479351.
70. Freitas K, Carroll FI, Damaj MI (2013) The antinociceptive effects of nicotinic receptors $\alpha 7$ -positive allosteric modulators in murine acute and tonic pain models. *J Pharmacol Exp Ther* 344: 264-275. PubMed: 23115222.
71. Freitas K, Negus SS, Carroll FI, Damaj MI (2013) In vivo pharmacological interactions between a type II positive allosteric modulator of $\alpha 7$ nicotinic ACh receptors and nicotinic agonists in a murine tonic pain model. *Br J Pharmacol* 169: 567-579. doi:10.1111/j.1476-5381.2012.02226.x. PubMed: 23004024.
72. McLean SL, Grayson B, Idris NF, Lesage AS, Pemberton DJ et al. (2011) Activation of $\alpha 7$ nicotinic receptors improves phencyclidine-induced deficits in cognitive tasks in rats: Implications for therapy of cognitive dysfunction in schizophrenia. *Eur Neuropsychopharmacol* 21: 333-343. doi:10.1016/j.euroneuro.2010.06.003. PubMed: 20630711.
73. McLean SL, Idris NF, Grayson B, Gendle DF, Mackie C et al. (2012) PNU-120596, a positive allosteric modulator of $\alpha 7$ nicotinic acetylcholine receptors, reverses a sub-chronic phencyclidine-induced cognitive deficit in the attentional set-shifting task in female rats. *J Psychopharmacol* 26: 1265-1270. doi:10.1177/0269881111431747. PubMed: 22182741.
74. Jin K, Minami M, Lan JQ, Mao XO, Bateur S et al. (2001) Neurogenesis in dentate subgranular zone and rostral subventricular

- zone after focal cerebral ischemia in the rat. *Proc Natl Acad Sci U S A* 98: 4710-4715. doi:10.1073/pnas.081011098. PubMed: 11296300.
75. Bederson JB, Pitts LH, Germano SM, Nishimura MC, Davis RL et al. (1986) Evaluation of 2,3,5-triphenyltetrazolium chloride as a stain for detection and quantification of experimental cerebral infarction in rats. *Stroke* 17: 1304-1308. doi:10.1161/01.STR.17.6.1304. PubMed: 2433817.
 76. Swanson RA, Morton MT, Tsao-Wu G, Savalos RA, Davidson C et al. (1990) A semiautomated method for measuring brain infarct volume. *J Cereb Blood Flow Metab* 10: 290-293. doi:10.1038/jcbfm.1990.47. PubMed: 1689322.
 77. Bederson JB, Pitts LH, Tsuji M, Nishimura MC, Davis RL et al. (1986) Rat middle cerebral artery occlusion: evaluation of the model and development of a neurologic examination. *Stroke* 17: 472-476. doi:10.1161/01.STR.17.3.472. PubMed: 3715945.
 78. Sun F, Xie L, Mao X, Hill J, Greenberg DA et al. (2012) Effect of a contralateral lesion on neurological recovery from stroke in rats. *Restor Neurol Neurosci* 30: 491-495. PubMed: 22868223.
 79. DeMuth JE. *Basic Statistics and Pharmaceutical Statistical Applications*. New York, N.Y.: Marcel Dekker Inc./CRC Press.
 80. Fujiki M, Kobayashi H, Uchida S, Inoue R, Ishii K (2005) Neuroprotective effect of donepezil, a nicotinic acetylcholine-receptor activator, on cerebral infarction in rats. *Brain Res* 1043: 236-241. doi:10.1016/j.brainres.2005.02.063. PubMed: 15862539.
 81. Sitzia F, Brown JT, Randall AD, Dunlop J (2011) Voltage- and Temperature-Dependent Allosteric Modulation of $\alpha 7$ Nicotinic Receptors by PNU120596. *Front Pharmacol* 2: 81. PubMed: 22207849.
 82. Shimohama S (2009) Nicotinic receptor-mediated neuroprotection in neurodegenerative disease models. *Biol Pharm Bull* 32: 332-336. doi:10.1248/bpb.32.332. PubMed: 19252273.
 83. Olinicy A, Stevens KE (2007) Treating schizophrenia symptoms with an $\alpha 7$ nicotinic agonist, from mice to men. *Biochem Pharmacol* 74: 1192-1201. doi:10.1016/j.bcp.2007.07.015. PubMed: 17714692.
 84. Kem WR (2000) The brain $\alpha 7$ nicotinic receptor may be an important therapeutic target for the treatment of Alzheimer's disease: studies with DMXBA (GTS-21). *Behav Brain Res* 113: 169-181. doi:10.1016/S0166-4328(00)00211-4. PubMed: 10942043.
 85. Klein J, Holler T, Cappel E, Köppen A, Löffelholz K (1993) Release of choline from rat brain under hypoxia: contribution from phospholipase A2 but not from phospholipase D. *Brain Res* 630: 337-340. doi:10.1016/0006-8993(93)90674-C. PubMed: 8118702.
 86. Klein J, Köppen A, Löffelholz K (1998) Regulation of free choline in rat brain: dietary and pharmacological manipulations. *Neurochem Int* 32: 479-485. doi:10.1016/S0197-0186(97)00127-7. PubMed: 9676747.
 87. Uteshev VV, Meyer EM, Papke RL (2003) Regulation of neuronal function by choline and 4OH-GTS-21 through $\alpha 7$ nicotinic receptors. *J Neurophysiol* 89: 1797-1806. PubMed: 12611953.
 88. Papke RL, Porter Papke JK (2002) Comparative pharmacology of rat and human $\alpha 7$ nAChR conducted with net charge analysis. *Br J Pharmacol* 137: 49-61. doi:10.1038/sj.bjp.0704833. PubMed: 12183330.
 89. Rao AM, Hatcher JF, Dempsey RJ (2000) Lipid alterations in transient forebrain ischemia: possible new mechanisms of CDP-choline neuroprotection. *J Neurochem* 75: 2528-2535. PubMed: 11080206.
 90. Gasull T, DeGregorio-Rocasolano N, Zapata A, Trullas R (2000) Choline release and inhibition of phosphatidylcholine synthesis precede excitotoxic neuronal death but not neurotoxicity induced by serum deprivation. *J Biol Chem* 275: 18350-18357. doi:10.1074/jbc.M910468199. PubMed: 10748226.
 91. Djuricic B, Olson SR, Assaf HM, Whittingham TS, Lust WD et al. (1991) Formation of free choline in brain tissue during in vitro energy deprivation. *J Cereb Blood Flow Metab* 11: 308-313. doi:10.1038/jcbfm.1991.63. PubMed: 1997502.
 92. Kiewert C, Mdzinarishvili A, Hartmann J, Bickel U, Klein J (2010) Metabolic and transmitter changes in core and penumbra after middle cerebral artery occlusion in mice. *Brain Res* 1312: 101-107. doi:10.1016/j.brainres.2009.11.068. PubMed: 19961839.
 93. Kihara T, Shimohama S, Sawada H, Honda K, Nakamizo T et al. (2001) $\alpha 7$ nicotinic receptor transduces signals to phosphatidylinositol 3-kinase to block A β amyloid-induced neurotoxicity. *J Biol Chem* 276: 13541-13546. PubMed: 11278378.
 94. Akaïke A, Takada-Takatori Y, Kume T, Izumi Y (2010) Mechanisms of neuroprotective effects of nicotine and acetylcholinesterase inhibitors: role of $\alpha 4$ and $\alpha 7$ receptors in neuroprotection. *J Mol Neurosci* 40: 211-216. doi:10.1007/s12031-009-9236-1. PubMed: 19714494.
 95. Shaw S, Bencherif M, Marrero MB (2002) Janus kinase 2, an early target of $\alpha 7$ nicotinic acetylcholine receptor-mediated neuroprotection against A β (1-42) amyloid. *J Biol Chem* 277: 44920-44924. doi:10.1074/jbc.M204610200. PubMed: 12244045.
 96. Hardingham GE, Bading H (2003) The Yin and Yang of NMDA receptor signalling. *Trends Neurosci* 26: 81-89. doi:10.1016/S0166-2236(02)00040-1. PubMed: 12536131.
 97. Papadia S, Hardingham GE (2007) The dichotomy of NMDA receptor signaling. *Neuroscientist* 13: 572-579. doi:10.1177/1073858407305833. PubMed: 18000068.
 98. Collins F, Schmidt MF, Guthrie PB, Kater SB (1991) Sustained increase in intracellular calcium promotes neuronal survival. *J Neurosci* 11: 2582-2587. PubMed: 1714495.
 99. Koike T, Martin DP, Johnson EM Jr. (1989) Role of Ca²⁺ channels in the ability of membrane depolarization to prevent neuronal death induced by trophic-factor deprivation: evidence that levels of internal Ca²⁺ determine nerve growth factor dependence of sympathetic ganglion cells. *Proc Natl Acad Sci U S A* 86: 6421-6425. doi:10.1073/pnas.86.16.6421. PubMed: 2548215.
 100. Blair LA, Bence-Hanulec KK, Mehta S, Franke T, Kaplan D et al. (1999) Akt-dependent potentiation of L channels by insulin-like growth factor-1 is required for neuronal survival. *J Neurosci* 19: 1940-1951. PubMed: 10066247.
 101. Del Barrio L, Martín-de-Saavedra MD, Romero A, Parada E, Egea J et al. (2011) Neurotoxicity induced by okadaic acid in the human neuroblastoma SH-SY5Y line can be differentially prevented by $\alpha 7$ and $\beta 2^*$ nicotinic stimulation. *Toxicol Sci* 123: 193-205. doi:10.1093/toxsci/kfr163. PubMed: 21715663.
 102. Castro NG, Albuquerque EX (1995) α -Bungarotoxin-sensitive hippocampal nicotinic receptor channel has a high calcium permeability. *Biophys J* 68: 516-524. doi:10.1016/S0006-3495(95)80213-4. PubMed: 7696505.
 103. Uteshev VV (2010) Evaluation of Ca²⁺ permeability of nicotinic acetylcholine receptors in hypothalamic histaminergic neurons. *Acta Biochim Biophys Sin (Shanghai)* 42: 8-20. doi:10.1093/abbs/gmp101. PubMed: 20043042.
 104. Fucile S (2004) Ca²⁺ permeability of nicotinic acetylcholine receptors. *Cell Calcium* 35: 1-8. doi:10.1016/j.ceca.2003.08.006. PubMed: 14670366.
 105. Bok J, Wang Q, Huang J, Green SH (2007) CaMKII and CaMKIV mediate distinct pro-survival signaling pathways in response to depolarization in neurons. *Mol Cell Neurosci* 36: 13-26. doi:10.1016/j.mcn.2007.05.008. PubMed: 17651987.
 106. Vaillant AR, Mazzoni I, Tudan C, Boudreau M, Kaplan DR et al. (1999) Depolarization and neurotrophins converge on the phosphatidylinositol 3-kinase-Akt pathway to synergistically regulate neuronal survival. *J Cell Biol* 146: 955-966. doi:10.1083/jcb.146.5.955. PubMed: 10477751.
 107. Si ML, Lee TJ (2001) Presynaptic $\alpha 7$ -nicotinic acetylcholine receptors mediate nicotine-induced nitric oxidergic neurogenic vasodilation in porcine basilar arteries. *J Pharmacol Exp Ther* 298: 122-128. PubMed: 11408533.
 108. Si ML, Lee TJ (2002) $\alpha 7$ -nicotinic acetylcholine receptors on cerebral perivascular sympathetic nerves mediate choline-induced nitric neurogenic vasodilation. *Circ Res* 91: 62-69. doi:10.1161/01.RES.0000024417.79275.23. PubMed: 12114323.
 109. Parada E, Egea J, Buendia I, Negrodo P, Cunha AC et al. (2013) The Microglial $\alpha 7$ -Acetylcholine Nicotinic Receptor Is a Key Element in Promoting Neuroprotection by Inducing Heme Oxygenase-1 via Nuclear Factor Erythroid-2-Related Factor. p. 2. *Antioxid Redox Signal*.
 110. Sharma G, Vijayaraghavan S (2001) Nicotinic cholinergic signaling in hippocampal astrocytes involves calcium-induced calcium release from intracellular stores. *Proc Natl Acad Sci U S A* 98: 4148-4153. doi:10.1073/pnas.071540198. PubMed: 11259680.
 111. Shytle RD, Mori T, Townsend K, Vendrame M, Sun N et al. (2004) Cholinergic modulation of microglial activation by $\alpha 7$ nicotinic receptors. *J Neurochem* 89: 337-343. doi:10.1046/j.1471-4159.2004.02347.x. PubMed: 15056277.
 112. De Rosa MJ, Dionisio L, Agriello E, Bouzat C, Esandi Mdel C (2009) $\alpha 7$ nicotinic acetylcholine receptor modulates lymphocyte activation. *Life Sci* 85: 444-449. doi:10.1016/j.lfs.2009.07.010. PubMed: 19632243.
 113. Wang H, Yu M, Ochani M, Amella CA, Tanovic M et al. (2003) Nicotinic acetylcholine receptor $\alpha 7$ subunit is an essential regulator of inflammation. *Nature* 421: 384-388. doi:10.1038/nature01339. PubMed: 12508119.
 114. Munro G, Hansen RR, Erichsen HK, Timmermann DB, Christensen JK et al. (2012) $\alpha 7$ nicotinic ACh receptor agonist compound B and positive allosteric modulator PNU-120596 both alleviate inflammatory hyperalgesia and cytokine release in the rat. *Br J Pharmacol*.
 115. Callahan PM, Hutchings EJ, Kille NJ, Chapman JM, Terry AV Jr. (2013) Positive allosteric modulator of $\alpha 7$ nicotinic-acetylcholine receptors, PNU-120596 augments the effects of donepezil on learning and memory in aged rodents and non-human primates.

- Neuropharmacology 67: 201-212. doi:10.1016/j.neuropharm.2012.10.019. PubMed: 23168113.
116. Chilton M, Mastropaolo J, Rosse RB, Bellack AS, Deutsch SI (2004) Behavioral consequences of methyllycaconitine in mice: a model of alpha7 nicotinic acetylcholine receptor deficiency. *Life Sci* 74: 3133-3139. doi:10.1016/j.lfs.2003.11.012. PubMed: 15081578.
117. Kelso ML, Pauly JR (2011) Therapeutic targets for neuroprotection and/or enhancement of functional recovery following traumatic brain injury. *Prog Mol Biol. Transl Sci* 98: 85-131.